

Award Number: DAMD17-98-1-8058

TITLE: Genetics of PTEN in Cowden Syndrome and Sporadic Breast Cancer

PRINCIPAL INVESTIGATOR: Charis Eng, M.D., Ph.D.

CONTRACTING ORGANIZATION: The Ohio State University Research Foundation  
Columbus, Ohio 43210-1239

REPORT DATE: October 1999

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;  
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20001115 068

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204 Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 1999	3. REPORT TYPE AND DATES COVERED Annual (01 Oct 98 - 30 Sep 99)	
4. TITLE AND SUBTITLE Genetics of PTEN in Cowden Syndrome and Sporadic Breast Cancer			5. FUNDING NUMBERS DAMD17-98-1-8058	
6. AUTHOR(S) Charis Eng, M.D., Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Ohio State University Research Foundation Columbus, Ohio 43210-1239  e-mail: eng-l@medctr.osu.edu			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words)  Cowden syndrome (CS) is an autosomal dominant disorder characterized by multiple hamartomas and a high risk of breast, thyroid and other cancers. The susceptibility gene is PTEN. The edict of this grant is to determine the genetic role of PTEN in non-CS CS-like families or individuals. The PI has accrued 15 site specific breast cancer families, without known BRCA1 and BRCA2 mutations. No intragenic PTEN mutations were found in these families. To date, 70 CS-like families and individuals have been accrued. One occult germline PTEN intragenic mutation was found among these families. The mutation positive family has breast, thyroid and endometrial cancers. Unfortunately, only 5 other families have endometrial carcinoma. In summary, at least 1.5% of non-CS CS-like families carry occult PTEN mutations. This has implications for the proband and family with respect to cancer risk and surveillance. Our preliminary data may suggest that endometrial cancer is a true component of CS and the data suggests that the International Cowden Consortium clinical diagnostic criteria are robust. Further accrual of these CS-like families, enriching for endometrial cancer, will be achieved and PTEN analysis, including the promoter, pursued.				
14. SUBJECT TERMS Breast Cancer			15. NUMBER OF PAGES 54	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

## FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

\_\_\_\_\_ Where copyrighted material is quoted, permission has been obtained to use such material.

\_\_\_\_\_ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

\_\_\_\_\_ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

\_\_\_\_\_ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

☒ \_\_\_\_\_ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

\_\_\_\_\_ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

\_\_\_\_\_ In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

\_\_\_\_\_ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.



10.1.99

\_\_\_\_\_  
PI - Signature

\_\_\_\_\_  
Date

## Table of Contents

Front Cover	1
SF 298	2
Foreword	3
Table of Contents	4
Introduction	5
Body	6
Key Research Accomplishments	7
Reportable Outcomes	7
Conclusions	8
References	9
Appendix	11

### Introduction

Cowden syndrome (CS) is an autosomal dominant disorder characterized by multiple hamartomas and a high risk of breast, thyroid and other cancers (reviewed by Eng <sup>1</sup>). The risk of breast cancer in affected women can range from 25-50% <sup>2,3</sup>. The PI mapped the CS gene to 10q22-23 <sup>4</sup> and subsequently identified *PTEN*, encoding a dual specificity phosphatase, as the or at least a major CS susceptibility gene <sup>5</sup>. That *PTEN* is a major CS gene was subsequently confirmed by other groups <sup>6-8</sup>. In addition, the PI has shown that germline *PTEN* mutations cause a proportion of Bannayan-Riley-Ruvalcaba syndrome (BRR), an autosomal dominant disorder characterized by megencephaly, mental retardation, lipomatosis, and speckled penis, previously thought not to be associated with cancer <sup>9</sup>.

Because CS is difficult to diagnose and is under-recognized and therefore under-diagnosed, the PI chairing the International Cowden Consortium synthesized a set of diagnostic criteria for the operational diagnosis of CS (Table 1) <sup>10</sup>, initially for research purposes and now, for clinical diagnostic purposes as well.

Table 1. International Cowden Consortium Diagnostic Criteria for CS

#### Pathognomonic Criteria

Mucocutaneous lesions:

- Trichilemmomas, facial
- Acral keratoses
- Papillomatous papules
- Mucosal lesions

#### Major Criteria

- Breast CA
- Thyroid CA, esp. follicular thyroid carcinoma
- Macrocephaly (Megalencephaly) (say,  $\geq 97\%$ ile)
- Lhermitte-Duclos disease (LDD)

#### Minor Criteria

- Other thyroid lesions (e.g adenoma or multinodular goiter)
- Mental retardation (say,  $IQ \leq 75$ )
- GI hamartomas
- Fibrocystic disease of the breast
- Lipomas
- Fibromas
- GU tumors (eg uterine fibroids) or malformation

#### Operational Diagnosis in an Individual:

1. Mucocutaneous lesions alone if:
  - a) there are 6 or more facial papules, of which 3 or more must be trichilemmoma, or
  - b) cutaneous facial papules and oral mucosal papillomatosis, or
  - c) oral mucosal papillomatosis and acral keratoses, or

- d) palmo plantar keratoses, 6 or more
- 2. 2 Major criteria but one must include macrocephaly or LDD
- 3. 1 Major and 3 minor criteria
- 4. 4 minor criteria

Operational Diagnosis in a Family where One Individual is Diagnostic for Cowden

- 1. The pathognomonic criterion/ia
- 2. Any one major criterion with or without minor criteria
- 3. Two minor criteria

The question of "cryptic" CS or the frequency of mutation-proven CS in individuals or families presenting with components of CS, such as breast cancer and/or thyroid cancer and/or endometrial cancer is important for the patient and his/her family with regard to medical management. In this regard, this proposal asks two main questions:

Task 1: What proportion of familial breast cancer only families have CS?

Task 2: What proportion of CS-like families, which do not make the full diagnostic criteria of the International Cowden Consortium (Table 1) have germline *PTEN* mutations, with its full implications, targeting cases and families that have breast and/or thyroid/endometrial cancer.

**Body**

**Task 1: Germline *PTEN* mutations in breast cancer only families**

To date, a total of 15 site specific breast cancer families that are mutation negative for *BRCA1* and *BRCA2* have been accrued and documented. All 15 are germline *PTEN* mutation negative. Approximately half of the affected tested individuals are heterozygous at *PTEN* IVS8+32T/G, thus excluding whole gene deletion. At this point, there is extensive data to show that gross gene deletion only results in BRR and even then, it is rare <sup>11</sup>. Again, all this analysis has been performed from small amounts of DNA left from old samples used for *BRCA* searches or from paraffin-embedded archived material, thus making Southern analysis impossible on these particular samples.

The plans for Years 2 and 3 is to continue accrual of site specific breast cancer families without *BRCA1* and *BRCA2* mutations for PCR-based germline *PTEN* analysis. These hopefully will be from peripheral blood leucocytes, thus making Southern analysis possible. The promoter lies within or is 250 kb long. Efforts by our lab and other labs are beginning to determine which portion or all of this segment is the minimal "true" promoter. If feasible after these analyses, then promoter mutation analysis will be performed in these families as well.

It would also appear from recent data (from our lab and others) that *PTEN* can be silenced or "inactivated" by causes other than structural gene alteration (ie mutation or deletion) (eg, Dahia et al 1999 <sup>12</sup>). In this regard, the PI plans to extend Task 1 to include examination of *PTEN* expression using immunohistochemistry to delineate

whether structural and/or other epigenetic phenomena pertain in PTEN inactivation in breast carcinogenesis.

**Task 2: Mutation Analysis in Non-CS Breast-Thyroid and/or Endometrial Carcinoma Families/Individuals ("CS-Like Families")**

To date, a total of 70 individuals or families with a CS-like syndrome have been accrued and cancers and tumors in affected individuals have been documented either with pathology report (preferable), death certificate or physician's notes. Each of these cases or families does not meet the operational diagnostic criteria for CS (Table 1). Further, they must minimally have at least one member with nonmedullary thyroid carcinoma and at least one other related member with breast cancer diagnosed at any age. They could also comprise single cases with both nonmedullary thyroid tumor and breast cancer. Among these 70 families/individuals, 1 germline *PTEN* mutation, c.209T->C (exon 3), was detected<sup>13</sup> (unpublished data). This was detected in a family where the proband was diagnosed with follicular thyroid carcinoma at the age of 31 and his mother had breast carcinoma diagnosed at 49 and 53, respectively, and endometrial carcinoma at 63. Half of these families/individuals were heterozygous at IVS8+32T/G thus excluding whole gene deletion. We and others have also shown that in CS and even BRR, whole gene deletion is rare<sup>5,7,9,11,14,15</sup>. Indeed, if *PTEN* is grossly deleted, only the BRR phenotype results<sup>11</sup>. Therefore, for the moment, the PI has decided that further hemizygote analysis on the large scale is not cost-efficient nor scientifically warranted.

From this Year 1 analysis, it would appear that the endometrial cancer feature in CS-like cases and families might increase the likelihood of finding a germline *PTEN* mutation. Therefore, while accrual of further CS-like families will continue, the PI will target families with endometrial pathology for Years 2 and 3. Southern analysis must await further accrual as amount of DNA per sample is limited with much of the analyses of these first 70 occurring off DNA templates extracted from paraffin-embedded archival material. Although promoter analysis was proposed in the original SOW, the promoter is within a 250 kb segment. We and others are trying to determine if the true promoter may be within a more confined region before beginning genetic analyses – this strategy would be meaningful.

### Key Research Accomplishments

#### Task 1

- Clinical-genetic database being set up

#### Task 2

- Clinical-genetic database of CS-like families being set up
- Delineate the frequency of occult germline *PTEN* mutation in CS-like families and individuals
- Discovered that based on molecular data, the clinical operational diagnostic criteria for classic CS is robust
- Based on the findings thus far, the PI as Chair of the International Cowden Consortium has recommended that endometrial carcinoma be added to the list of major criteria in the operational diagnosis of CS (see Table 1 for 1995 Criteria). This is the first data-based update of criteria since 1995. The US NCCN/Genetic-High Risk Panel has agreed to adopt this revision in their 2000 guidelines.

### Reportable Outcomes

Marsh DJ, Dahia PLM, Caron S, Kum JB, Frayling IM, Tomlinson IPM, Hughes KS, Hodgson SV, Murday VA, Houlston R, **Eng C**. Germline *PTEN* mutations in Cowden syndrome-like families. J Med Genet 1998; 35:881-5.



### Conclusions

In Year 1 of the grant, the PI has continued to accrue non-CS families and individuals. A clinical-genetic database is actively being built. It is envisioned that this will be on-going for the next 2 years. Because of the PI's disruptive move from Boston in the beginning of the year, the database assistant is just being hired (the PI herself was manually inputting and analyzing data). Nonetheless, in the first analysis of non-CS CS-like families and individuals, the PI has found approximately 1.5-2% with an occult germline *PTEN* mutation. Re-examination of the family has found that they have breast, thyroid and endometrial cancer and no other stigmata of CS. Thus, the PI preliminarily concludes that the Clinical Operational Criteria for CS Diagnosis proposed by the International Cowden Consortium is robust, and that perhaps, endometrial carcinoma should be added to the list of major criteria.

In order to confirm these early findings, the PI will continue to accrue such CS-like families and individuals but to enrich for endometrial cancer or non-neoplastic endometrial disease (eg young onset endometrial fibroids can be a feature of CS). Germline *PTEN* mutation analysis will be pursued and the promoter elucidated and finally examined for alterations.

## References

1. Eng, C. Genetics of Cowden syndrome - through the looking glass of oncology. *Intl. J. Oncol.* 1998; **12**:701-710.
2. Starink, TM, van der Veen, JPW, Arwert, F, de Waal, LP, de Lange, GG, Gille, JJP, Eriksson, AW. The cowden syndrome: a clinical and genetic study in 21 patients. *Clin Genet* 1986; **29**:222-233.
3. Longy, M, Lacombe, D. Cowden disease. Report of a family and review. *Ann. Génét.* 1996; **39**:35-42.
4. Nelen, MR, Padberg, GW, Peeters, EAJ, Lin, AY, van den Helm, B, Frants, RR, Coulon, V, Goldstein, AM, van Reen, MMM, Easton, DF, Eeles, RA, Hodgson, S, Mulvihill, JJ, Murday, VA, Tucker, MA, Mariman, ECM, Starink, TM, Ponder, BAJ, Ropers, HH, Kremer, H, Longy, M, Eng, C. Localization of the gene for Cowden disease to 10q22-23. *Nature Genet.* 1996; **13**:114-116.
5. Liaw, D, Marsh, DJ, Li, J, Dahia, PLM, Wang, SI, Zheng, Z, Bose, S, Call, KM, Tsou, HC, Peacocke, M, Eng, C, Parsons, R. Germline mutations of the *PTEN* gene in Cowden disease, an inherited breast and thyroid cancer syndrome. *Nature Genet.* 1997; **16**:64-67.
6. Lynch, ED, Ostermeyer, EA, Lee, MK, Arena, JF, Ji, H, Dann, J, Swisshelm, K, Suchard, D, MacLeod, PM, Kvinnsland, S, Gjertsen, BT, Heimdal, K, Lubs, H, Moller, P, King, M-C. Inherited mutations in *PTEN* that are associated with breast cancer, Cowden syndrome and juvenile polyposis. *Am. J. Hum. Genet.* 1997; **61**:1254-1260.
7. Tsou, HC, Teng, D, Ping, XL, Broncolini, V, Davis, T, Hu, R, Xie, X-X, Gruener, AC, Schrager, CA, Christiano, AM, Eng, C, Steck, P, Ott, J, Tavtigian, SV, Peacocke, M. Role of *MMAC1* mutations in early onset breast cancer: causative in association with Cowden's syndrome and excluded in *BRCA1*-negative cases. *Am. J. Hum. Genet.* 1997; **61**:1036-1043.
8. Nelen, MR, van Staveren, CG, Peeters, EAJ, Ben Hassel, M, Gorlin, RJ, Hamm, H, Lindboe, CF, Fryns, J-P, Sijmons, RH, Woods, DG, Mariman, ECM, Padberg, GW, Kremer, H. Germline mutations in the *PTEN/MMAC1* gene in patients with Cowden disease. *Hum. Mol. Genet.* 1997; **6**:1383-1387.
9. Marsh, DJ, Dahia, PLM, Zheng, Z, Liaw, D, Parsons, R, Gorlin, RJ, Eng, C. Germline mutations in *PTEN* are present in Bannayan-Zonana syndrome. *Nature Genet.* 1997; **16**:333-334.
10. Eng, C. Cowden syndrome. *J. Genet. Counsel.* 1997; **6**:181-191.
11. Marsh, DJ, Kum, JB, Lunetta, KL, Bennett, MJ, Gorlin, RJ, Ahmed, SF, Bodurtha, J, Crowe, C, Curtis, MA, Dazouki, M, Dunn, T, Feit, H, Geraghty, MT, Graham, JM, Hodgson, SV, Hunter, A, Korf, BR, Manchester, D, Miesfeldt, S, Murday, VA, Nathanson, KA, Parisi, M, Poher, B, Romano, C, Tolmie, JL, Trembath, R, Winter, RM, Zackai, EH, Zori, RT, Weng, LP, Dahia, PLM, Eng, C. *PTEN* mutation spectrum and genotype-phenotype correlations in Bannayan-Riley-Ruvalcaba syndrome suggest a single entity with Cowden syndrome. *Hum. Mol. Genet.* 1999; **8**:1461-1472.
12. Dahia, PLM, Aguiar, RCT, Alberta, J, Kum, J, Caron, S, Sills, H, Marsh, DJM, Freedman, A, Ritz, J, Stiles, C, Eng, C. *PTEN* is inversely correlated with the cell survival factor PKB/Akt and is inactivated by diverse mechanisms in haematologic malignancies. *Hum. Mol. Genet.* 1999; **8**:185-193.

13. Marsh, DJ, Caron, S, Dahia, PLM, Kum, JB, Frayling, IM, Tomlinson, IPM, Hughes, KS, Hodgson, SV, Murday, VA, Houlston, R, Eng, C. Germline *PTEN* mutations in Cowden syndrome-like families. *J. Med. Genet.* 1998; **35**:881-885.
14. Marsh, DJ, Coulon, V, Lunetta, KL, Rocca-Serra, P, Dahia, PLM, Zheng, Z, Liaw, D, Caron, S, Duboué, B, Lin, AY, Richardson, A-L, Bonnetblanc, J-M, Bressieux, J-M, Cabarrot-Moreau, A, Chompret, A, Demange, L, Eeles, RA, Yahanda, AM, Fearon, ER, Fricker, J-P, Gorlin, RJ, Hodgson, SV, Huson, S, Lacombe, D, LePrat, F, Odent, S, Toulouse, C, Olopade, OI, Sobol, H, Tishler, S, Woods, CG, Robinson, BG, Weber, HC, Parsons, R, Peacocke, M, Longy, M, Eng, C. Mutation spectrum and genotype-phenotype analyses in Cowden disease and Bannayan-Zonana syndrome, two hamartoma syndromes with germline *PTEN* mutation. *Hum. Mol. Genet.* 1998; **7**:507-515.
15. Tsou, HC, Ping, XL, Xie, XX, Gruener, AC, Zhang, H, Nini, R, Swisshelm, K, Sybert, V, Diamond, TM, Sutphen, R, Peacocke, M. The genetic basis of Cowden's syndrome: three novel mutations in PTEN/MMAC1/TEP1. *Hum. Genet.* 1998; **102**:467-473.

## Appendix

Marsh DJ, Dahia PLM, Caron S, Kum JB, Frayling IM, Tomlinson IPM, Hughes KS, Hodgson SV, Murday VA, Houlston R, **Eng C**. Germline *PTEN* mutations in Cowden syndrome-like families. J Med Genet 1998; 35:881-5.

## Original articles

# Germline PTEN mutations in Cowden syndrome-like families

Debbie J Marsh, Patricia L M Dahia, Stacey Caron, Jennifer B Kum, Ian M Frayling, Ian P M Tomlinson, Kevin S Hughes, Rosalind A Eeles, Shirley V Hodgson, Vicky A Murday, Richard Houlston, Charis Eng

Department of Adult Oncology and Charles A Dana Human Cancer Genetics Unit, Dana-Farber Cancer Institute, Department of Medicine, Harvard Medical School, Susan and Richard Smith Laboratories, SM822, 1 Jimmy Fund Way, Boston, MA 02115-6084, USA  
D J Marsh  
P L M Dahia  
S Caron  
J B Kum  
C Eng

ICRF Colorectal Cancer Unit, St Mark's Hospital, Northwick Park, Middlesex, UK  
I M Frayling

Tumour Genetics Group, Wellcome Trust Centre for Human Genetics, University of Oxford, UK  
I P M Tomlinson

Department of Surgery, Lahey Clinic, Burlington, MA 02115, USA  
K S Hughes

Section of Molecular Carcinogenesis, Institute of Cancer Research and Cancer Genetics Clinic, Royal Marsden Hospital, Sutton, Surrey, UK  
R A Eeles  
R Houlston

Department of Medical Genetics, United Medical and Dental Schools of Guy's and St Thomas's Hospital, London, UK  
S V Hodgson

Department of Clinical Genetics, St George's Hospital Medical School, London, UK  
V A Murday

## Abstract

Cowden syndrome (CS) or multiple hamartoma syndrome (MIM 158350) is an autosomal dominant disorder with an increased risk for breast and thyroid carcinoma. The diagnosis of CS, as operationally defined by the International Cowden Consortium, is made when a patient, or family, has a combination of pathognomonic major and/or minor criteria. The CS gene has recently been identified as PTEN, which maps at 10q23.3 and encodes a dual specificity phosphatase. PTEN appears to function as a tumour suppressor in CS, with between 13-80% of CS families harbouring germline nonsense, missense, and frameshift mutations predicted to disrupt normal PTEN function. To date, only a small number of tumour suppressor genes, including BRCA1, BRCA2, and p53, have been associated with familial breast or breast/ovarian cancer families. Given the involvement of PTEN in CS, we postulated that PTEN was a likely candidate to play a role in families with a "CS-like" phenotype, but not classical CS. To answer these questions, we gathered a series of patients from families who had features reminiscent of CS but did not meet the Consortium Criteria. Using a combination of denaturing gradient gel electrophoresis (DGGE), temporal temperature gel electrophoresis (TTGE), and sequence analysis, we screened 64 unrelated CS-like subjects for germline mutations in PTEN. A single male with follicular thyroid carcinoma from one of these 64 (2%) CS-like families harboured a germline point mutation, c.209T→C. This mutation occurred at the last nucleotide of exon 3 and within a region homologous to the cytoskeletal proteins tensin and auxilin. We conclude that germline PTEN mutations play a relatively minor role in CS-like families. In addition, our data would suggest that, for the most part, the strict International Cowden Consortium operational diagnostic criteria for CS are quite robust and should remain in place.

(J Med Genet 1998;35:881-885)

Keywords: PTEN; Cowden syndrome; breast; thyroid

Breast and thyroid carcinoma are two frequently occurring neoplasms in the female population. Increased risks for both breast and thyroid cancer are prominent features of Cowden syndrome (CS). The hallmark phenotype of this inherited cancer syndrome is the presence of hamartomas, developmentally incorrect, benign, hyperplastic growths, in multiple organ systems including the skin, gastrointestinal tract, central nervous system, breast, and thyroid. Breast cancer will develop in 25-50% of women with CS and 3-10% of all CS patients will develop thyroid cancer.<sup>1,2</sup> At present, only four tumour suppressor genes have been associated with familial breast cancer, BRCA1, BRCA2, p53, and PTEN.<sup>3-7</sup> Initially thought to account for over 80% of hereditary breast cancer,<sup>8,9</sup> germline mutations in BRCA1 and BRCA2 together are now thought to account for 25-50% of all familial breast cancer,<sup>10</sup> thus opening up the possibility of other BRCAX genes. Along these lines, germline mutations in p53 are associated with 70% of cases of Li-Fraumeni syndrome, an autosomal dominant condition comprising breast cancer, brain tumours, sarcomas, and adrenocortical carcinomas.<sup>3,4,11</sup> Recently, the CS susceptibility gene has been identified as the tumour suppressor gene PTEN, also known as MMAC1 and TEP1.<sup>7,12-14</sup> PTEN maps to 10q23.3 and encodes a 403 amino acid dual specificity phosphatase.<sup>12-15</sup> Germline missense and truncating mutations have been reported in between 13-80% of patients with CS.<sup>7,16-18</sup> It should be noted that while initial linkage studies of 12 families with CS was highly suggestive of a single locus for CS,<sup>19</sup> a subsequent study proposes that genetic heterogeneity may exist in CS.<sup>16</sup>

At the somatic level, PTEN has been shown to be mutated or deleted in a number of human malignancies, including sporadic breast, brain, prostate, and kidney cancer cell lines, as well as in a number of primary tumours including endometrial carcinomas, glioblastomas, malignant melanoma, and thyroid and breast tumours.<sup>20-33</sup>

Given the role of PTEN in CS and the relatively large percentage of familial cases of breast cancer that are not caused by germline mutation of BRCA1, BRCA2, or p53, we sought to determine whether PTEN may be mutated in

Table 1 Phenotypic classification of CS-like families

Phenotype of families	No of families
Breast and thyroid carcinoma occurring together in at least one person	22
Breast and thyroid carcinoma occurring in different subjects	32
Breast carcinoma and thyroid disease (eg goitre)	3
Breast carcinoma/CS-like (eg trichilemmoma), no thyroid involvement	6
Thyroid carcinoma/CS-like, no breast involvement	1
Total	64




Figure 1 DGGE detection of *c.209T→C* in the germline of a patient from a CS-like family. Control mutations from CS and BRR families are also included to display the sensitivity of this technique for the detection of PTEN mutations. Lane 1, wild type control (exon 3); lane 2, Y68H (exon 3); lane 3, IVS2-2A→G (exon 3); lane 4, *c.209T→C* (exon 3); lane 5, wild type control (amplicon 5I, representing the 5' half of exon 5); lane 6, Q87X (amplicon 5I); lane 7, *c.347-351delACAAT* (amplicon 5I); lane 8, wild type control (amplicon 5II, representing the 3' half of exon 5); lane 9, C124R (amplicon 5II); lane 10, E157X (amplicon 5II); lane 11, wild type control (exon 7); lane 12, R233X (exon 7); lane 13, *c.791ATins* (exon 7).

the germline of families that did not meet the strict diagnostic criteria for CS determined by the International Cowden Consortium.<sup>2,19</sup> The phenotypes of these families were, minimally, breast and non-medullary thyroid cancers, and, maximally, a sum of phenotypes falling just short of the Consortium Criteria for CS.

### Material and methods

#### PATIENTS

Members of 64 unrelated CS-like families were collected for analysis (table 1). These CS-like families were defined as families or people that have some, but not all, of the features of CS and do not meet the operational diagnostic criteria of the International Cowden Consortium. Minimally, these CS-like families contained at least one member with both non-medullary thyroid cancer and at least one other related member with breast cancer diagnosed at any age. They also could comprise subjects with both breast cancer and non-medullary thyroid cancer. Alternatively, families could be made up of either breast or non-medullary thyroid cancer and other features of CS, such as trichilemmomas, without meeting the consortium criteria for CS.

The diagnostic criteria for classical CS used in this study has been previously described by the Consortium.<sup>2,19</sup> In brief, the diagnosis of CS requires that a patient or family meet a combination of pathognomonic major and minor criteria. Major criteria include breast cancer, non-medullary thyroid cancer (especially follicular thyroid carcinoma), macrocephaly ( $\geq 97$ th centile), and Lhermitte-Duclos disease (LDD), which is a dysplastic gangliocytoma of the cerebellum that can cause seizures, tremors, and poor coordination. Hamartomas of

the skin, including trichilemmomas (benign tumours of the hair follicle infundibulum) and mucocutaneous papillomatous papules (for example, scrotal tongue), are diagnostic if there are six or more papules, with three or more being trichilemmomas. Minor criteria include benign thyroid lesions such as multinodular goitre and adenomas, fibrocystic breast disease, mental retardation ( $IQ \leq 75$ ), gastrointestinal hamartomas, lipomas, fibromas, and genitourinary tumours or malformations. Individual people or families would be diagnosed with CS if they have two major criteria, where one is either LDD or macrocephaly, one major with three minor criteria, or four minor criteria. No patients in this study fulfilled these criteria. Constitutional DNA was extracted from blood leucocytes using standard, previously described methods.<sup>34</sup> Approval for the use of human subjects in this study was obtained under IRB approved protocol 94-138 (Dana-Farber Cancer Institute).

#### DENATURING GRADIENT GEL ELECTROPHORESIS (DGGE) AND TEMPORAL TEMPERATURE GEL ELECTROPHORESIS (TTGE)

A combination of DGGE and TTGE was performed for all nine exons of PTEN. GC clamped primer sequences, PCR conditions, and DGGE conditions have been previously described,<sup>35</sup> with the exception of primers for exons 2 and 4. Exon 2 and 4 primer sequences, with GC clamps added, were as follows: exon 2, 2F, 5'-CGT CCC GCG TTT GAT TGC TGC ATA TTT CAG-3' and 2R, 5'-CGC CCG CCG CGC CCC GCG CCC GTC CCG CCG CCC CCG CCC GTC TAA ATG AAA ACA CAA CAT G-3'; exon 4, 4F, 5'-CGC CCG CCG CGC CCC GCG CCC GTC CCG CCG CCC CCG CCC GAA ATA ATA AAC ATT ATA AAG ATT CAG GCA ATG-3' and 4R, 5'-GAC AGT AAG ATA CAG TCT ATC-3'. Split exon 5 primers with GC clamps and conditions for mutation detection have been previously reported.<sup>26</sup>

TTGE is a mutation detection technique using the basic PCR fragment denaturation principles of DGGE. The major difference between these methods is that a temperature gradient, rather than a chemical gradient of varying urea and glycerol percentages, is used for strand separation of the GC clamped homo- and heteroduplexed PCR products by generating a linear temperature gradient over the length of the electrophoresis run (Bio-Rad Laboratories, Hercules, CA). One or 0.75 mm thick gels of 10% polyacrylamide:bis (37.5:1) (Bio-Rad Laboratories) and 7 mol/l urea (Bio-Rad Laboratories) were run using the DCode™ Universal Mutation Detection System (Bio-Rad Laboratories). Electrophoresis was performed at 130 V for six hours with a temperature gradient of 46-58°C and a ramp rate of 2°C per hour. TTGE fragments were visualised under ultraviolet transillumination after the gel was stained with ethidium bromide (Bio-Rad Laboratories).

Both DGGE and TTGE have proven high accuracy in detecting mutations in general and specifically in detecting known PTEN mutations from CS patients (fig 1).

CRC Human Cancer Genetics Research Group, University of Cambridge, Addenbrooke's Hospital, Box 238, Level 3, Laboratories Block, Hills Road, Cambridge CB2 2QQ, UK  
C Eng

Correspondence to: Dr Eng, Human Cancer Genetics Program, Ohio State University Comprehensive Cancer Center, 420 W 12th Avenue, 690 MRF, Columbus, OH 43210, USA.

Received 19 March 1998  
Revised version accepted for publication 23 April 1998

## SEQUENCE ANALYSIS

Exons which showed DGGE and TTGE variants underwent direct sequence analysis. The PCR primers and reaction conditions have been described elsewhere.<sup>7 14 28 35</sup> PCR products were gel isolated and purified using the Wizard PCR Preps DNA Purification System (Promega, Madison, WI). Direct sequencing of these products was performed using the ABI Prism dye terminator cycle sequencing ready reaction kit (Perkin-Elmer Corp. Norwalk, CT). Cycle sequencing products were electrophoresed on 6% Long ranger gels (FMC Bioproducts, Rockland, ME) and analysed on an Applied Biosystems model 373A automated DNA sequencer (Perkin-Elmer Corp).

## PTEN POLYMORPHISM ANALYSIS

A previously identified intronic polymorphic site in PTEN, IVS8+32G/T, was analysed in a single affected member from each CS-like family to investigate hemizyosity at the PTEN locus in mutation negative families. This site is moderately heterozygous, with an earlier report finding 50% of samples to be informative.<sup>28</sup> Potential hemizyosity was assessed by the amplification of exon 8 and flanking intronic sequence and digestion with the restriction endonuclease *HincII* under conditions suggested by the manufacturer (New England Biolabs, Beverly, MA).

## Results

## PTEN MUTATION ANALYSIS

A missense point mutation, c.209T→C (L70P), predicted to affect splicing was identified in a single affected patient (1 of 64, 2%) (fig 1). This mutation was not identified in 100 normal alleles. When this occult germline PTEN mutation was identified, the family history was reassessed (fig 2). The subject analysed for this study, III.1, developed follicular thyroid carcinoma at the age of 31. His

mother, II.2, had breast adenocarcinoma diagnosed at the age of 49 and again at 53. She also had endometrial carcinoma diagnosed at 63 years. Careful clinical assessment of these two subjects was unable to identify macrocephaly, skin lesions typical of CS, or scrotal tongue. The maternal grandfather, I.1, was diagnosed with leukaemia at the age of 57. Unfortunately, family members other than III.1 were unavailable for analysis. Fresh tumour from III.1, which would have allowed us to study the putative aberrant splicing effect of this mutation, was also unavailable. No mutations were identified in the other 63 unrelated CS-like families.

## PTEN POLYMORPHISM ANALYSIS

Forty-eight percent (30 of 63) of unrelated subjects from PTEN mutation negative CS-like families were found to be heterozygous at the IVS8+32T/G site. This analysis would suggest that, at least in these families, gross germline deletion of PTEN can be excluded.

## Discussion

An occult germline PTEN mutation, c.209T→C at the last nucleotide of exon 3 was found in one of 64 (2%) CS-like families. This family's cancers, comprising leukaemia, which may or may not be related, adenocarcinoma of the breast, endometrial carcinoma, and follicular thyroid carcinoma, together do not meet the International Cowden Consortium Criteria used for the diagnosis of CS in this study. However, we cannot exclude the possibility that this family represents a case of low penetrance CS. The family with PTEN mutation in this study contrasts with that in a recent study that reported a PTEN mutation in a family initially classified as having breast and thyroid tumours only but reclassified as CS after mutation analysis led to closer clinical assessment.<sup>36</sup> Closer clinical assessment of the family presented in the current study did not identify additional features of CS.

In the remaining families where no occult germline mutations were identified, it is highly unlikely that these mutations would have gone undetected. Both DGGE and TTGE are highly sensitive mutation detection techniques<sup>37</sup> and both have been shown consistently to detect known PTEN mutations and other sequence polymorphisms (Marsh and Eng, unpublished data, 1998; fig 1). Further, because at least one affected member from nearly half of these mutation negative families was heterozygous at the IVS8+32T/G polymorphism, whole gene deletion is unlikely, at least in these families.

In CS, while missense and truncating mutations are scattered largely along the entirety of PTEN, a mutational "hot spot" exists in exon 5, which contains the PTPase core motif at codons 122-132.<sup>7 16-18</sup> Thus, many mutations in CS are predicted to disrupt the phosphatase function of this protein. Interestingly, the mutation identified in exon 3 falls in the N-terminal half of the PTEN protein that has been shown to have some sequence similarities to the cytoskeletal proteins tensin and auxilin.




Figure 2 Pedigree of CS-like family with the occult germline PTEN mutation. c.209T→C was identified in DNA extracted from blood leucocytes from patient III.1 who presented with follicular thyroid carcinoma.

Specifically, the leucine residue at codon 70 that is altered by this T to C point mutation (L70P) is conserved in both bovine auxilin and chicken tensin.<sup>14</sup> Thus, it is possible that this mutation may be affecting the phosphatase function of this protein, as one may predict if this putative splice site mutation leads to a truncated protein, and may also function to disrupt normal cellular motility and cell-cell interactions.

Whether germline PTEN mutations are associated with CS and related inherited hamartoma syndromes (Bannayan-Ruvalcaba-Riley syndrome, (BRR, MIM 153480) and juvenile polyposis syndrome (JPS, MIM 174900)), as well as syndromes comprising partial CS phenotypes, is largely unknown. Before the identification of PTEN as the CS gene, it was not inconceivable that the three related hamartoma syndromes and CS-like syndromes were all associated with different mutations in a single gene. We have shown that germline PTEN mutations are associated with the great majority, approximately 80%, of classical CS families.<sup>7,18</sup> Nelen *et al*<sup>17</sup> identified PTEN mutations in 47% of CS cases studied. One other study of 23 CS families identified only 13% of families with germline PTEN mutation.<sup>16</sup> This was perhaps not surprising as limited linkage information in these families suggested the possibility of genetic heterogeneity in CS, even though initial studies of a group of 12 CS families showed no evidence for heterogeneity.<sup>19</sup>

We have also shown that germline PTEN mutations account for at least a proportion of BRR, which is characterised by macrocephaly, lipomatosis, thyroid dysfunction, hamartomatous polyps of the gastrointestinal tract, and pigmented macules of the glans penis, but without a known predisposition to breast and thyroid cancer.<sup>18,38</sup> How mutations in a single gene, at times identical,<sup>18,38</sup> can function to predispose to two overlapping but apparently distinct syndromes, one with malignancy and one without, remains to be elucidated.

Disparate reports concerning the third hamartoma syndrome, JPS, and PTEN mutation or deletion have recently been published.<sup>35,36,39-41</sup> A putative JPS locus, JP1, at 10q22-24 was initially thought to encompass PTEN, although fine structure mapping placed this locus slightly centromeric of PTEN.<sup>42</sup> Subsequently, the 10q22-24 region was excluded as a putative JPS locus by linkage analysis in eight JPS families.<sup>35</sup> Screening of PTEN in 21 classical JPS families and 16 cases of sporadic JPS did not identify any germline mutations.<sup>35,39</sup> In contrast, PTEN mutation has been reported in four patients with "juvenile polyposis",<sup>36,41</sup> although the clinical diagnosis of classic juvenile polyposis in these cases is questionable. Given these genetic data and the phenotypic overlap of these syndromes, we can say with some confidence that if a germline PTEN mutation were detected in a person previously thought to have "juvenile polyposis", then the diagnosis needs to be revised, as that person is likely to have either CS or BRR.

Along the same lines, we have now investigated a cohort of families, each of which contains some of the component tumours of CS but do not meet the Consortium diagnostic criteria for CS. Only one such family was found to have an occult germline PTEN mutation, arguing that such germline alterations play a minor role in families that do not meet the strict CS diagnostic criteria. Nonetheless, this finding is significant for three reasons. Firstly, it suggests that the operational diagnostic criteria for CS established by the International Cowden Consortium are, for the most part, robust and are useful for identifying PTEN mutation positive CS families. Secondly, we must also conclude from our data that other genes are involved which lend susceptibility to a CS-like disease and to site specific breast and non-medullary thyroid cancer. Thirdly, for non-CS subjects identified with occult PTEN mutations, albeit uncommonly, there are important implications for future hamartoma/cancer development that should impact on surveillance.

Unanswered questions remain, however. For example, are CS-like families without germline PTEN mutations at any less risk of cancer than those with mutations? Preliminary genotype-phenotype analyses suggest that classical CS families without germline PTEN mutations are at lower risk of developing malignant breast disease compared to their PTEN mutation positive counterparts.<sup>18</sup> By extrapolation, it would seem that PTEN mutation negative CS-like families should be at decreased risk of developing breast cancer. Unfortunately, this study was unable to confirm this clinically relevant extrapolation. We can conclude, however, that in the majority of cases, germline PTEN mutations lead specifically to a CS or BRR phenotype and that the phenotype of CS-like families is, for the most part, caused by unknown mechanisms.

We would like to thank the patients and families who participated in this study, and Dr Oliver Gimm for critical reading of this manuscript. Ms Elaine Krekonis is acknowledged for assistance with the patients. The Molecular Biology Core Facility at the Dana-Farber Cancer Institute, Boston is acknowledged for running sequencing gels. This study was supported by the Susan G Komen Breast Cancer Foundation (to PLMD and CE), the American Cancer Society (RPG 97-064-02VM), the Barr Investigatorship, and a Breast Cancer Research Grant (34088PP1009) from the Massachusetts Department of Public Health (to CE). CE is the Lawrence and Susan Marx Investigator in Human Cancer Genetics.

- 1 Starink TM, van der Veen JPW, Arwert F, *et al*. The Cowden syndrome: a clinical and genetic study in 21 patients. *Clin Genet* 1986;29:222-33.
- 2 Eng C. Cowden syndrome. *J Genet Counsel* 1997;6:9181-92.
- 3 Malkin D, Li FP, Fraumeni JF Jr, *et al*. Germ-line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms. *Science* 1990;250:1233-8.
- 4 Srivastava S, Zou Z, Pirolo K, Blattner W, Chang EH. Germ-line transmission of a mutated p53 gene in a cancer-prone family with Li-Fraumeni syndrome. *Nature* 1990;348:747-9.
- 5 Miki Y, Swensen J, Shattuck-Eidens D, *et al*. A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. *Science* 1994;266:66-71.
- 6 Wooster R, Bignell G, Lancaster J, *et al*. Identification of the breast cancer susceptibility gene BRCA2. *Nature* 1995;378:789-92.
- 7 Liaw D, Marsh DJ, Li J, *et al*. Germline mutations of the PTEN gene in Cowden disease, an inherited breast and thyroid cancer syndrome. *Nat Genet* 1997;16:64-7.
- 8 Easton DF, Bishop DT, Ford D, Crockford GP, Consortium BCL. Genetic linkage analysis in familial breast and ovarian cancer: results from 214 families. The Breast Cancer Linkage Consortium. *Am J Hum Genet* 1993;52:678-701.



- 9 Wooster R, Neuhausen SL, Manjion J, *et al.* Localization of a breast cancer susceptibility gene, *BRCA2*, to chromosome 13q12-12. *Science* 1994;265:2088-90.
- 10 Eng C. From bench to bedside...but when? *Genome Res* 1997;7:669-72.
- 11 Varley JM, McGown G, Thorncroft M, *et al.* Germ-line mutations of TP53 in Li-Fraumeni families: an extended study of 39 families. *Cancer Res* 1997;57:3245-52.
- 12 Li J, Yen C, Liaw D, *et al.* *PTEN*, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. *Science* 1997;275:1943-7.
- 13 Li DM, Sun H. TEP1, encoded by a candidate tumor suppressor locus, is a novel protein tyrosine phosphatase regulated by transforming growth factor  $\beta$ . *Cancer Res* 1997;57:2124-9.
- 14 Steck PA, Pershouse MA, Jasser SA, *et al.* Identification of a candidate tumor suppressor gene, *MMAC1*, at chromosome 10q23.3 that is mutated in multiple advanced cancers. *Nat Genet* 1997;15:356-62.
- 15 Myers MP, Stolarov JP, Eng C, *et al.* P-TEN, the tumor suppressor from human chromosome 10q23, is a dual-specificity phosphatase. *Proc Natl Acad Sci USA* 1997;94:9052-7.
- 16 Tsou HC, Teng DHF, Li Ping X, *et al.* The role of *MMAC1* mutations in early-onset breast cancer: causative in association with Cowden syndrome and excluded in *BRCA1*-negative cases. *Am J Hum Genet* 1997;61:1036-43.
- 17 Nelen MR, van Staveren WCG, Peeters EAJ, *et al.* Germline mutations in the *PTEN/MMAC1* gene in patients with Cowden disease. *Hum Mol Genet* 1997;6:1383-7.
- 18 Marsh DJ, Coulon V, Lunetta KL, *et al.* Mutation spectrum and genotype-phenotype analyses in Cowden disease and Bannayan-Zonana syndrome, 2 hamartoma syndromes with germline *PTEN* mutation. *Hum Mol Genet* 1998;7:507-15.
- 19 Nelen MR, Padberg GW, Peeters EAJ, *et al.* Localization of the gene for Cowden disease to chromosome 10q22-23. *Nat Genet* 1996;13:114-16.
- 20 Liu W, James CD, Frederick L, Alderete BE, Jenkins RB. *PTEN/MMAC1* mutations and EGFR amplification in glioblastomas. *Cancer Res* 1997;57:5254-7.
- 21 Teng DHF, Hu R, Lin H, *et al.* *MMAC1/PTEN* mutations in primary tumor specimens and tumor cell lines. *Cancer Res* 1997;57:5221-5.
- 22 Rhei E, Kang L, Bogomolny F, Federici MG, Borgen PJ, Boyd J. Mutation analysis of the putative tumor suppressor gene *PTEN/MMAC1* in primary breast carcinomas. *Cancer Res* 1997;57:3657-9.
- 23 Wang SI, Puc J, Li J, *et al.* Somatic mutations of *PTEN* in glioblastoma multiforme. *Cancer Res* 1997;57:4183-6.
- 24 Ahmed Rasheed BK, Stenzel TT, McLendon RE, *et al.* *PTEN* gene mutations are seen in high-grade but not in low-grade gliomas. *Cancer Res* 1997;57:4187-90.
- 25 Cairns P, Ookami K, Halachmi S, *et al.* Frequent inactivation of *PTEN/MMAC1* in primary prostate cancer. *Cancer Res* 1997;57:4997-5000.
- 26 Guldberg P, Straten PT, Birck A, Ahrenkiel V, Kirkin AF, Zeuthen J. Disruption of the *MMAC1/PTEN* gene by deletion or mutation is a frequent event in malignant melanoma. *Cancer Res* 1997;57:3660-3.
- 27 Tashiro H, Blazes MS, Wu R, *et al.* Mutations in *PTEN* are frequent in endometrial carcinoma but rare in other common gynecological malignancies. *Cancer Res* 1997;57:3935-40.
- 28 Dahia PLM, Marsh DJ, Zheng Z, *et al.* Somatic deletions and mutations in the Cowden disease gene, *PTEN*, in sporadic thyroid tumors. *Cancer Res* 1997;57:4710-13.
- 29 Risinger JJ, Hayes AK, Berchuk A, Barrett JC. *PTEN/MMAC1* mutations in endometrial cancers. *Cancer Res* 1997;57:4736-8.
- 30 Funari FB, Lin H, Su Huang HJ, Cacence WK. Growth suppression of glioma cells by *PTEN* requires a functional phosphatase catalytic domain. *Proc Natl Acad Sci USA* 1997;94:12479-84.
- 31 Sakurada A, Suzuki A, Sato M, *et al.* Infrequent genetic alterations of the *PTEN/MMAC1* gene in Japanese patients with primary cancers of the breast, lung, pancreas, kidney and ovary. *Jpn J Cancer Res* 1997;88:1025-8.
- 32 Ueda K, Nishijima M, Inui H, *et al.* Infrequent mutations in the *PTEN/MMAC1* gene among primary breast cancers. *Jpn J Cancer Res* 1997;89:17-21.
- 33 Chiariello E, Roz L, Albarosa R, Magnani I, Finocchiaro J. *PTEN/MMAC1* mutations in primary glioblastomas and short-term cultures of malignant gliomas. *Oncogene* 1998;16:541-5.
- 34 Mathew CG, Smith BA, Thorpe K, *et al.* Deletion of genes on chromosome 1 in endocrine neoplasia. *Nature* 1987;328:524-6.
- 35 Marsh DJ, Roth S, Lunetta KL, *et al.* Exclusion of *PTEN* and 10q22-24 as the susceptibility locus for juvenile polyposis syndrome. *Cancer Res* 1997;57:5017-21.
- 36 Lynch ED, Ostermeyer EA, Lee MK, *et al.* Inherited mutations in *PTEN* that are associated with breast cancer, Cowden disease, and juvenile polyposis. *Am J Hum Genet* 1997;61:1254-60.
- 37 Eng C, Vijg J. Genetic testing: the problems and the promise. *Nat Biotechnol* 1997;15:422-6.
- 38 Marsh DJ, Dahia PLM, Zheng Z, *et al.* Germline mutations in *PTEN* are present in Bannayan-Zonana syndrome. *Nat Genet* 1997;16:333-4.
- 39 Riggins GJ, Hamilton SR, Kinzler KW, Vogelstein B. Normal *PTEN* gene in juvenile polyposis. *J Neg Obstet Genet Oncol* 1998;1:1.
- 40 Tsuchiya KD, Wiesner G, Cassidy SB, Limwongse C, Boyle JT, Schwartz S. Deletion 10q23.2-q23.33 in a patient with gastrointestinal juvenile polyposis and other features of a Cowden-like syndrome. *Genes Chrom Cancer* 1997;21:113-18.
- 41 Olschwang S, Serova-Sinilnikova OM, Lenoir GM, Thomas G. *PTEN* germline mutations in juvenile polyposis coli. *Nat Genet* 1998;18:12-14.
- 42 Jacoby RF, Schlack S, Cole CE, Skarbek M, Harris C, Meisner LF. A juvenile polyposis tumor suppressor locus at 10q22 is deleted from nonepithelial cells in the lamina propria. *Gastroenterology* 1997;112:1398-403.

## CURRICULUM VITAE

Date Prepared: Sept 29, 1999

**Name:** Charis Eu Li ENG

**Office Address:** Human Cancer Genetics Program, Comprehensive Cancer Center,  
Ohio State University, 690C Medical Research Facility, 420 West  
12<sup>th</sup> Avenue, Columbus, OH 43210, USA

**Home Address:** 1683 Quarry Trace, Columbus, OH 43204, USA

**Place of Birth:** Republic of Singapore

### Education:

1982	BA	University of Chicago, Chicago, IL (Biological Sciences with Honors)
1986	PhD	University of Chicago, Chicago, IL (Developmental Biology)
1988	MD	University of Chicago, Chicago, IL (Medicine)

### Postdoctoral Training:

#### Internship and Residency:

1988-89	Intern, Internal Medicine, Beth Israel Hospital, Boston, MA
1989-90	Junior Assistant Resident, Internal Medicine, Beth Israel Hospital, Boston, MA
1990-91	Senior Assistant Resident, Internal Medicine, Beth Israel Hospital, Boston, MA

#### Fellowships:

1988-93	Clinical Fellow in Medicine, Harvard Medical School, Boston, MA
1991-94	Clinical Fellow, Division of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA
1991-92	Clinical Fellow, Division of Medical Oncology, Brigham and Women's Hospital, Boston, MA
1992-95	Cancer Research Campaign - Dana-Farber Cancer Institute Fellow in Human Cancer Genetics, University of Cambridge, UK
1992-95	Research Fellow, Department of Pathology, University of Cambridge, UK

### Licensure and Certification:

1990	Commonwealth of Massachusetts Medical Licensure, No. 72073
1991	American Board of Internal Medicine, Specialty Board Certification in General Internal Medicine, No. 135435
1992-95	Limited Registration, No. 92/3382, General Medical Council, London, UK
1997	American Board of Internal Medicine, Subspecialty Board Certification in Medical Oncology
119.	State of Ohio Medical Licensure

### Academic Appointments

1994-95	Instructor in Medicine, Harvard Medical School, Boston, MA
1995-98	Assistant Professor of Medicine, Harvard Medical School, Boston, MA
1999-	Associate Professor of Medicine and Human Cancer Genetics, Ohio State University, Columbus, OH

### Hospital or Affiliated Institution Appointments:

1992-95	Honorary Clinical Fellow/Senior Registrar, Clinical Cancer Genetics, Department of Clinical Genetics, Addenbrooke's Hospital, Cambridge, UK
1993-94	Honorary Clinical Status in Clinical Cancer Genetics, The Royal Marsden Hospital, London and Sutton, UK
1994-95	Honorary Consultant in Clinical Cancer Genetics, The Royal Marsden Hospital, London and Sutton, UK
1995-98	Active Staff Physician, Department of Adult Oncology, Dana-Farber Cancer Institute, Boston, MA
1995-98	Associate Physician, Division of Medical Oncology, Department of Medicine, Brigham and Women's Hospital, Boston, MA
1999-	Director, Clinical Cancer Genetics Program, James Cancer Hospital and Solove Research Institute, Comprehensive Cancer Center, Ohio State University, Columbus, OH
1999-	Member, Molecular Biology and Cancer Genetics Program, Comprehensive Cancer Center, Ohio State University, Columbus, OH

### Other Professional Positions and Major Visiting Appointments

1994-95	Member, Emmanuel College, Cambridge, UK
1995-	Honorary Member, Emmanuel College, Cambridge, UK
1995	Consultant to the Molecular Genetics Laboratory, Albert Ludwigs-Universität Freiburg, Abteilung Innere Medizin IV - Nephrologie, Freiburg, Germany, July 17-19
1995-	Honorary Fellow, CRC Human Cancer Genetics Research Group, University of Cambridge, UK

### Hospital and Health Care Organisation Clinical Responsibilities:

1995-98	Staff Medical Oncologist, Gastrointestinal Cancer Center, Dana-Farber Cancer Institute, Boston, MA
1995-98	Staff Clinical Cancer Geneticist, Cancer Risk and Prevention Clinic, Dana-Farber Cancer Institute, Boston, MA
1995-96	Staff Medical Oncologist, Head and Neck Clinic, Dana-Farber Cancer Institute, Boston, MA
1997-98	Staff Medical Oncologist, Endocrine Cancer Clinic, Dana-Farber Partners Cancer Center, Boston, MA
1999-	Director and Attending Clinical Cancer Geneticist, Clinical Cancer Genetics Program, James Cancer Hospital and Solove Research Institute, Comprehensive Cancer Center, Ohio State University, Columbus, OH

**Major Administrative Responsibilities:**

1996-99	Coordinator, Harvard Longwood Seminars in the Genetics of Cancer and Aging, Dana-Farber Partners Cancer Center, Boston, MA
1999-	Director, Clinical Cancer Genetics Program, James Cancer Hospital and Solove Research Institute, Comprehensive Cancer Center, Ohio State University, Columbus, OH

## Major Committee Assignments:

### Medical School

- 1983-86 Interviewer for first year applicants to the Pritzker School of Medicine, University of Chicago, IL  
 1999- Alternate Member, Biomedical Human Protection Committee, Ohio State University, Columbus, OH

### Hospital

- 1995 Scientific Steering Subcommittee, Gastrointestinal Cancer Center, Dana-Farber Cancer Institute, Boston, MA  
 1996-98 Molecular Diagnostics Committee of the Clinical Cancer Genetics Program, Dana-Farber Partners Cancer Care, Boston, MA  
 1996 High Risk Committee, Gastrointestinal Cancer Center, Dana-Farber Partners Cancer Care, Boston, MA  
 1996-98 Steering Committee, Endocrine Cancer Clinic, Dana-Farber Partners Cancer Center, Boston, MA  
 1997-98 Steering Committee, Gastrointestinal Cancer Center, Dana-Farber Partners Cancer Center, Boston, MA  
 1997-98 Human Cancer Genetics Working Group, Dana-Farber Partners Cancer Center, Boston, MA  
 1999- Clinical Trials Office Steering Committee, James Cancer Hospital and Solove Research Institute and Comprehensive Cancer Center, Ohio State University, Columbus, OH  
 1999- Clinical Scientific Research Committee, James Cancer Hospital and Solove Research Institute, Comprehensive Cancer Center, Ohio State University, Columbus, OH

### National

- 1996- Reviewer, Department of Veterans Affairs Merit Review Applications  
 1997-98 Reviewer and Expert Consultant, American Society of Clinical Oncology Task Force on Cancer Genetics Education  
 1998- Reviewer, Molecular Biology 3 Study Section, Department of Defence US Army Research Medical and Material Command Breast Cancer Research Program  
 1999 Reviewer, Susan G. Komen Breast Cancer Research Foundation Grants  
 1999 Site Visit Team Member, Quadriannual Site Visit, National Institute of Child Health and Development, Developmental Endocrinology Branch  
 1999 Reviewer, Cancer Genetics Section, American Society of Human Genetics Annual Meeting Abstracts  
 1999- National Comprehensive Cancer Network (NCCN) Guidelines Panel Member: Genetics/Familial High Risk Screening Guidelines

### International

- 1994- Coordinator and co-chair, International *RET* Mutation Consortium  
 1994- Coordinator and chair, International Cowden Syndrome Consortium  
 1995-98 International Review Board, Dutch Cancer Society  
 1997 Ad Hoc Review Committee, Programme Project Grant, National Cancer Institute of Canada

- 1997- Peer Review Panel, Project Grants, Comitato Promotore Telethon, Italy
- 1997- Reviewer, Project Grants and Clinical Research Fellowships, Cancer Research Campaign, London, UK
- 1997- Reviewer and Full Member, National Cancer Institute of Canada, Panel J: Pathology, Tumor Markers, Molecular Epidemiology and Clinical Correlative Studies, Toronto, ON
- 1998- Ad Hoc External Reviewer, Italian Association for Cancer Research
- 1998- Member, Steering Committee, Breast Cancer Information Core (BIC)
- 1999- Member, International Scientific Committee, 8<sup>th</sup> International Workshop on Multiple Endocrine Neoplasia, Jerusalem, Israel, May 2001

### **Professional Societies and Colleges:**

- 1982- Phi Beta Kappa, Member
- 1982-87 Sigma Xi, Associate Member
- 1982-88 American Medical Students' Association, Member
- 1982-88 American Medical Association, Member
- 1982-89 American Medical Women's Association, Member
- 1984-88 American Association for the Advancement of Science, Member
- 1984-89 New York Academy of Sciences, Member
- 1987- Sigma Xi, Member
- 1988- Alpha Omega Alpha, Member
- 1989-92 American College of Physicians, Associate
- 1990-98 Massachusetts Medical Society, Member
- 1992-99 American College of Physicians, Member
- 1995- New York Academy of Sciences, Member
- 1996- American Society of Clinical Oncology, Member
- 1996- American Society of Human Genetics, Member
- 1998- American Association for Cancer Research, Member
- 1999- American College of Physicians, Fellow

### **Editorial Boards:**

#### **Editorial Boards:**

- 1998- Journal of Medical Genetics, North American Editor
- 1998- Journal of Medical Genetics, Associate Editor for Cancer Genetics
- 1998- Journal of Endocrine Genetics, Editorial Board Member

#### ***Ad hoc* Reviewer for:**

- 1998- American Journal of Human Genetics
- 1997- American Journal of Pathology
- 1997- American Journal of Surgical Pathology
- 1999- BioTechniques
- 1997- Blood
- 1998- British Journal of Cancer
- 1998- Cancer
- 1996- Cancer Epidemiology, Biomarkers and Prevention
- 1997- Cancer Research

1997-	Carcinogenesis
1998-	Clinical Cancer Research
1995-	Clinical Endocrinology
1995-	Clinical Genetics
1996-	European Journal of Endocrinology
1997-	European Journal of Human Genetics
1997-	Experimental Cell Research
1996-	Gastroenterology
1995-	Genes, Chromosomes and Cancer
1998-	Genomics
1997-	Human Genetics
1994-	Human Molecular Genetics
1994-	Human Mutation
1998-	International Journal of Cancer
1996-	Journal of the American Medical Association
1995-	Journal of Clinical Endocrinology and Metabolism
1999-	Journal of Experimental Medicine
1999-	Journal of Clinical Investigation
1995-	Journal of Clinical Oncology
1994-98	Journal of Medical Genetics
1998-	Journal of the National Cancer Institute
1996	Mutation Research
1995-	Nature Genetics
1996-	New England Journal of Medicine
1995-	Oncogene
1997-	Proceedings of the National Academy of Sciences, USA

#### Awards and Honors:

1978-82	Dean's List, College, University of Chicago, IL
1981	Edmondson Summer Research Fellowship, University of Chicago, IL
1981-82	Yim Chan Merit Scholarship, University of Chicago, IL
1982	Graduation with Divisional and Collegiate Honors, University of Chicago, IL
1982	Phi Beta Kappa
1982	Sigma Xi, Associate Membership
1982	Sigma Xi Science Prize Competition, Honorable Mention, University of Chicago, IL
1982	Sigma Xi Certificate of Merit for Excellence in Undergraduate Scientific Research, University of Chicago, IL
1982-83	Dean's Letter of Commendation for Excellence in Gross Anatomy and Microbiology, Pritzker School of Medicine, University of Chicago, IL
1982-84	Far East Scholarship, Pritzker School of Medicine, University of Chicago, IL
1983	National Institutes of Health Summer Research Fellowship, Pritzker School of Medicine, University of Chicago, IL
1984-86	American Heart Association - Borg-Warner Medical Student Research Fellowship, University of Chicago, IL
1987	Sigma Xi, promotion to Full Membership
1988	Alpha Omega Alpha
1990	Nomination for Chief Residency, Department of Medicine, Beth Israel Hospital, Boston, for 1992-93 (Position Declined)

- 1990 Nomination for the National Institutes of Health-Upjohn Medical Residents Research Award
- 1991 Upjohn Travel Award to the Meeting of the American Association for Cancer Research, Houston, TX
- 1992 Johanna Wood Fellowship, Dana-Farber Cancer Institute, Boston, MA
- 1992-95 Cancer Research Campaign - Dana-Farber Cancer Institute Fellowship in Human Cancer Genetics, University of Cambridge, U.K.
- 1995-97 Lucille P. Markey Charitable Trust Young Scientist Award
- 1995-98 The First Lawrence and Susan Marx Investigatorship in Human Cancer Genetics, Dana-Farber Cancer Institute, Boston, MA
- 1996 Patterson Fellowship, Dana-Farber Cancer Institute, Boston, MA
- 1997-99 Barr Investigatorship, Dana-Farber Cancer Institute, Boston, MA
- 1999 International Scientific Committee, 8<sup>th</sup> International Workshop on Multiple Endocrine Neoplasia, Jerusalem, Israel, May, 2001



## Laboratory and Clinical Investigator Track

### A. Report of Research

#### 1. Major research interests:

1. Cancer Genetics
2. Molecular Epidemiology of Cancer
3. Second Malignancies in Retinoblastoma Patients
4. Genetics of Multiple Endocrine Neoplasia Type 2 and Related Cancers
5. Familial Gastrointestinal Cancers
6. Cowden Syndrome and Related Cancers
7. Inherited Hamartoma-Neoplasia Syndromes

#### 2. Narrative description of research

The broad thrust of my laboratory involves the utilisation of DNA-based methods to identify and characterise genes which cause susceptibility to inherited cancer syndromes, to determine their role in sporadic carcinogenesis and to perform molecular epidemiologic analyses as they might relate to future clinical applications. Upon this framework, we are examining the genetics of two inherited thyroid cancer syndromes, Cowden syndrome (nonmedullary thyroid cancer) and MEN 2 (medullary thyroid cancer), and related sporadic cancers. Hence, the genetics of susceptibility gene *PTEN*, encoding a dual specificity phosphatase on 10q23.3, is being examined in Cowden syndrome and other inherited hamartoma syndromes as well as populations of isolated breast and thyroid cancer cases. Somatic genetics of *PTEN* is being pursued in a range of sporadic cancers including sporadic counterpart Cowden component tumors, breast, thyroid and endometrial carcinomas. Gene-gene interactions and gene-environment interactions are beginning to be explored. Biochemical, cellular and functional studies are beginning to be performed in our laboratory as well as in collaboration with a number of laboratories locally, nationally and internationally. The genetics of the *RET* proto-oncogene are pursued for clinical translational purposes for MEN 2 and sporadic neuroendocrine tumors. Towards those ends, genotype-phenotype analyses and genotype-prognosis analyses are being performed. Examination of common low penetrance variants in sporadic medullary thyroid carcinoma is also being pursued in the hope of identifying common alleles for predisposition in sporadic neuroendocrine tumors.

Recent efforts in my laboratory have focused on the role of the nuclear receptor transcription factor PPAR $\gamma$  in sporadic carcinogenesis. Troglitazone (Rezulin<sup>TM</sup>), which is a specific synthetic ligand for PPAR $\gamma$ , is an oral hypoglycemic agent used by over 1.6 million Americans. So, our work may have broad implications not only for examining the pathogenesis of common cancers but may impact public health as well. This avenue of investigation also promises direct translation into clinical oncologic practice.

#### 3. Research funding information:

1981	Edmondson Summer Research Fellowship, University of Chicago (Advisor: Edward D. Garber)	PI
1978-82	Yim Chan Merit Scholarship, University of Chicago, IL	
1984-86	American Heart Association Borg-Warner Medical Student Research Fellowship, University of Chicago Pritzker School of Medicine, IL	PI
1992-95	Cancer Research Campaign [CRC] Dana-Farber Fellowship	PI

Integrated fellowship in clinical cancer genetics and molecular cancer genetics at the University of Cambridge, UK  
(Advisor: Bruce A. J. Ponder)

1995-97	New Investigator Award, Charles A. Dana Foundation	
1995-97	New Investigator Award, Markey Charitable Trust	
1995-98	Lawrence and Susan Marx Investigatorship in Human Cancer Genetics	PI
1996	Patterson Fellowship	PI
1996-98	Harvard Nathan Shock Center Award for the Basic Biology of Aging, NIA State of the art resource core for two dimensional gene scanning	
1996-99	Barr Investigatorship Human cancer genetics research	PI
1997-98	Women's Cancer Program Grant, Dana-Farber Partners Cancer Center Development of a rapid multi-gene test for hereditary breast cancer	PI
1997-99	American Cancer Society (National) Research Project Grant Isolation and characterisation of Cowden syndrome gene	PI
1997-1999	DFG Training Fellowship (Germany) Trainee PI: Oliver Gimm, MD Novel mutations and low penetrance alleles in the <i>RET</i> proto-oncogene in multiple endocrine neoplasia type 2 and sporadic medullary thyroid carcinoma	Mentor
1997-2000	Susan G. Komen Breast Cancer Foundation Postdoctoral Fellowship Trainee: Patricia L M Dahia, MD, PhD Role of Cowden susceptibility gene in breast cancer	PI
1998	Breast Cancer Research Award, Massachusetts Department of Public Health <i>PTEN</i> , the Cowden disease gene, in patients and families with breast cancer and thyroid disease	PI
1998-99	ASCO Young Investigator Award Prognostic markers for progression of esophageal adenocarcinoma Trainee PI: Matthew H. Kulke, MD	Mentor
1998-1999	Concert for the Cure Breast Cancer Research Award Genetics of <i>PTEN</i> in Cowden syndrome and unselected breast cancer patients	PI
1999	Ohio State University Seed Grant Mapping the susceptibility gene for hereditary and sporadic Barrett esophagus and esophageal adenocarcinoma	PI
1998-2001	Department of Defence US Army Breast Cancer Research Program Genetics of <i>PTEN</i> in different forms of hereditary breast cancer	PI
1998-2001	American Cancer Society (National) Research Project Grant Genetics of <i>PTEN</i> in Cowden syndrome and sporadic breast cancer	PI

- 1999-2002      National Institutes of Health Workstatement (RFP)  
A phase 2 study of a selective estrogen receptor modulator (LY353381) vs.  
Tamoxifen vs. placebo in premenopausal women with an increased risk for breast  
cancer
- 1999-2001      Mary Kay Ash Charitable Foundation Grant      PI  
Genetic and functional analysis of PPAR-gamma as a novel tumor suppressor locus  
in sporadic breast carcinoma

## B. Report of Teaching

### Local Contributions

#### Medical School / School of Public Health

- 1985 Medical Genetics, Teaching Assistant for 100-110 second year medical students, University of Chicago Pritzker School of Medicine (Contact 5 hr/wk, Prep 5 hr/wk)
- 1996-98 Molecular Epidemiology, Guest Lecturer for 30-50 medical, dental and graduate students, medical fellows and instructors, Harvard School of Public Health (Contact 1-2 hr, Prep 2 hr)
- 1997 HMS211A Graduate Course in Biochemistry and Cell Biology, invited lecture on inherited cancer syndromes for 20 graduate, dental and medical students, Harvard Medical School, Boston: (Contact 1.5 hr, Prep 2 hr)
- 1998 Harvard Medical School Course in Genetics, Embryology and Reproduction, Tutor for group of 7-10 medical students (Contact 40 hr, Prep 20 hr)

#### Graduate Medical Course/Seminar/Invited Teaching Presentation

- 1991 Grand Rounds, Beth Israel Hospital, Boston: Causes of late mortality in retinoblastoma patients, invited speaker (Contact 20 min, Prep 3 hr)
- 1994 Department of Medicine Seminar Series, University of Cambridge School of Clinical Medicine: The many faces of *RET*, invited lecture for 50 housestaff and faculty of the Clinical School (Contact 1 hr, Prep 2 hr)
- 1996 Seminars in Medicine of the Beth Israel Hospital: From bench to bedside: the *RET* proto-oncogene in multiple endocrine neoplasia, invited lecture for 30-60 faculty and trainees from the Boston area (Contact 1.5 hr, Prep 3 hr)
- 1996 Harvard Medical School Department of Genetics Seminar: The polygenic etiology of Hirschsprung disease, invited speaker for 20-25 clinical genetics fellows, postdoctoral fellows and genetics faculty (Contact 1 hr, Prep 2 hr)
- 1997 Brigham and Women's Hospital Specialty Lecture for Medical Housestaff: Genetics of endocrine tumors, invited speaker for 50-60 medical housestaff (Contact 1 hr, Prep 1 hr)
- 1997 Massachusetts Cancer Center Seminar, Charlestown, MA: *RET*, *GDNF* and *GDNFR- $\alpha$*  in MEN 2, invited speaker for 30-50 PIs, postdoctoral fellows and graduate students (Contact 1.5 hr, Prep 2 hr)
- 1997 GI Grand Rounds, Massachusetts General Hospital: Molecular genetics of Hirschsprung disease for 15-25 GI fellows and faculty (Contact 1 hr, Prep 2 hr)
- 1997 Women's Cancer Program, Dana-Farber Partners Cancer Center, Boston: Identification of the Cowden syndrome susceptibility gene, invited speaker for 20-30 multidisciplinary faculty, clinical fellows, housestaff, postdoctoral fellows, graduate students (Contact 1 hr, Prep 1 hr)
- 1997 Breast Center Basic Biology Seminar, Dana-Farber Partners Cancer Center, Boston: Identification of the Cowden syndrome gene, a multipurpose gene which predisposes to breast and thyroid cancers, invited speaker for 40-60 multidisciplinary faculty, fellows and housestaff (Contact 1 hr, Prep 1 hr)
- 1997 Harvard-Longwood Seminars in the Genetics of Cancer and Aging, Boston: *PTEN* in inherited hamartoma-cancer syndromes: one gene-many syndromes? Invited speaker for 50-70 clinical and basic science faculty, postdoctoral fellows, clinical fellows, and graduate students from the Harvard Longwood area (Contact 1 hr, Prep 1 hr)

- 1997 Massachusetts General Hospital Cancer Center Grand Rounds, Boston: *PTEN* in Cowden syndrome and sporadic breast and thyroid cancers (Contact 1 hr, Prep 1 hr)
- 1999 Ohio State University Human Cancer Genetics Program Seminar, Columbus, OH: *PTEN* and the great imitator: Cowden syndrome (Contact 1 hr, Prep 1 hr)

#### **Continuing Medical Education Course**

- 1997 Cancer Genetics for Office Practice: Genetics of thyroid cancer in everyday practice, faculty (Contact 3 hr, Prep 1 hr)
- 1997 American College of Surgeons, Massachusetts Chapter, Waltham: Genetics of colorectal tumors, faculty (Contact 2 hr, prep 1 hr)
- 1998 Massachusetts Eye and Ear Infirmary and Harvard Medical School Course on Thyroid and Parathyroid Tumors: *RET* and medullary thyroid carcinoma, faculty (Contact 30 min, prep 20 min)

#### **Advisory and Supervisory Responsibilities**

- 1988-89 Teaching and supervision of Harvard medical students during clinical clerkship, Beth Israel Hospital, 1 medical student per rotation (200 hr/yr)
- 1989-91 Teaching and supervision of Harvard medical students during clinical clerkship and medical interns, Beth Israel Hospital, 2-4 interns +/- 1 medical student per rotation (2000 hr/yr)
- 1991-92 Teaching and supervision of medical students, and medical housestaff from Brigham and Women's Hospital and Beth Israel Hospital, 3-8 housestaff +/- 1 medical student per month (500 hr/yr)
- 1993-95 Teaching and supervision of technicians, students and junior postdoctoral fellows, CRC Human Cancer Genetics Research Group, Department of Pathology, University of Cambridge, 2 technicians, 0-3 medical/graduate students and 0-1 junior postdoctoral fellow (20 hr/wk)
- 1995- Teaching and supervision of postdoctoral fellows, students and technicians working in my laboratory, 2-6 postdoctoral fellows, 0-1 medical students, 1-3 technicians (15 hr/wk)
- 1996-98 Teaching and supervision of medical oncology and genetics fellows and genetics counsellors, Cancer Risk and Prevention Clinic, Dana-Farber Cancer Institute (3-5 hr/wk)
- 1996-98 Clinic Attending for medical oncology fellows, Dana-Farber Cancer Institute, 1-6 fellows per session (5-10 hr/mth)
- 1999- Direction and administration of the Clinical Cancer Genetics Program, Comprehensive Cancer Center, Ohio State University: 1.5-2 MD attending clinical cancer geneticists, 0-1 oncology fellow, 0-1 medical resident, 3-4 cancer genetics counselors, 0-1 research assistant, 1 data manager and 2 executive support associates (20 hr/wk)

#### **Laboratory-Based Trainees**

##### Postdoctoral Trainees

Debbie J. Marsh, PhD 1996-99  
 Project: Genetics of Cowden syndrome and Bannayan-Riley-Ruvalcaba syndrome  
 Current Position: Lecturer, Dept of Medicine, University of Sydney School of Medicine, Sydney, Australia

Matthew H. Kulke, MD 1997-99

Project: Molecular epidemiology and prognostic markers in sporadic gastrointestinal cancers  
 Current Position: Instructor in Medicine, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA

Patricia L.M. Dahia, MD, PhD 1997-  
 Project: Somatic genetics and biochemical expression of *PTEN* in sporadic tumors  
 Current Position: Postdoctoral Senior Research Associate, Dana-Farber Cancer Institute and Harvard Medical School, Boston, MA; Instructor in Medicine, Harvard Medical School

Oliver Gimm, MD 1997-  
 Project: Genetics of neuroendocrine tumors  
 Current Position: DFG Postdoctoral Researcher, Human Cancer Genetics Program, Ohio State University, Columbus, OH

Aurel Perren, MD 1998  
 Project: Immunocytochemistry of *PTEN* in sporadic tumors of the breast and thyroid  
 Current Position: Resident in Pathology, University of Zürich School of Medicine, Zürich, Switzerland

Jen Jen Yeh, MD 1998-99  
 Project: Somatic genetics of non-medullary thyroid carcinomas and the role of the mitochondrial genome  
 Current Position: Postdoctoral Research Fellow, Dana-Farber Cancer Institute and Harvard Medical School, Boston, MA

Liang-Ping Weng, MD, MS 1998-  
 Project: Biochemistry and cell biology of *PTEN* in breast and thyroid carcinogenesis  
 Current Position: Research Scientist, Human Cancer Genetics Program, Ohio State University, Columbus, OH

Xiao-Ping Zhou, MD, PhD 1998-  
 Project: Genetics of central nervous system tumors  
 Current Position: Postdoctoral Researcher, Human Cancer Genetics Program, Ohio State University, Columbus, OH

Ravshan Burikhanov, PhD 1999-  
 Project: Cell biology of RET, *PTEN* and PPAR $\gamma$  in thyroid cancer models  
 Current Position: Postdoctoral Researcher, Human Cancer Genetics Program, Ohio State University, Columbus, OH

Keisuke Kurose, MD, PhD 1999-  
 Project: Genetics of *PTEN* and PPAR $\gamma$  in gynecologic cancers  
 Current Position: Postdoctoral Researcher, Human Cancer Genetics Program, Ohio State University Columbus, OH

#### Student Trainees

Antje Gössling 1996  
 Project: Genetics of GDNF and GFR $\alpha$ -1 in central nervous system tumors  
 Current Position: Resident in Clinical Genetics, Faculty of Medicine, University of Tübingen School of Medicine, Germany

Eva-Maria Dürr 1998  
 Project: Genetics of *CUL2* and *VBP-1* in pheochromocytomas  
 Current Position: Senior Medical Student, University of Bonn School of Medicine, Germany

Ying Huang 1999-  
 Project: Mapping the susceptibility gene for familial nonmedullary thyroid cancer  
 Role: PhD thesis committee member (Albert de la Chapelle, MD, PhD, Advisor and Chair)

### **Junior Faculty Mentored**

Matthew H. Kulke, MD Instructor in Medicine, Dana-Farber Cancer Institute  
 ASCO Young Investigator Award 1998-99

Kornelia Polyak, MD, PhD Assistant Professor of Medicine, Dana-Farber Cancer Institute  
 ASCO Career Development Award 1999-2003

Patricia L M Dahia, MD, PhD Instructor in Medicine, Dana-Farber Cancer Institute 1999-

Liang-Ping Weng, MD, MS Research Scientist, Ohio State University 1999-

### **Leadership Role**

1995-99 Director, Harvard Longwood Seminars in the Genetics of Cancer and Aging, organisation and coordination of seminar topic and speakers, invitation of speakers, and public relations for the seminar (CME 1 course)  
 1999- Director, Clinical Cancer Genetics Program, Comprehensive Cancer Center, Ohio State University

### Regional, National and International Contributions (Invited Presentations)

1993 Lancet Grand Round: Familial Cancer Syndromes.  
 Case Presentations and Multiple Endocrine Neoplasia Type 2A, Royal Marsden Hospital, Sutton  
 1993 ICRF Department of Medical Oncology Seminar, St. Bartholomew's Hospital, London: The multiple endocrine neoplasia type 2 syndromes  
 1994 Faculty, March of Dimes 25th Clinical Genetics Conference, Orlando, FL, USA Symposium in Genetics and Development: The molecular genetics of multiple endocrine neoplasia type 2  
 1994 Arbeitsgemeinschaft für Gynäkologische Onkologie, Vienna, Austria: The familial and genetic risks of ovarian cancer  
 1994 Postgraduate Training Course in Endocrinology: Multiple Endocrine Neoplasia Type 2. British Society for Endocrinology, St. Mary's Hospital, London, UK  
 1994 Symposium on Genotype-Phenotype Correlations, British Medical Genetics Conference, York, UK: Mutations of the *RET* proto-oncogene in the multiple endocrine neoplasia type 2 syndromes and Hirschsprung disease  
 1995 Case Presentation Conference, Department of Medical Genetics, BC Children's Hospital, University of British Columbia, Vancouver: The role of the *RET* proto-oncogene in the multiple endocrine neoplasia type 2 syndromes and Hirschsprung disease  
 1995 Meeting of the Clinical Molecular Genetics Society, Selwyn College, Cambridge: Mutational analysis of the *RET* proto-oncogene in MEN 2

- 1995 Department of Internal Medicine IV - Nephrology Special Seminar, Albert Ludwigs University of Freiburg, Germany: Pheochromocytoma and multiple endocrine neoplasia type 2: molecular genetic analysis
- 1995 EORTC Thyroid Group Meeting, London, UK: Germline mutations in the *RET* proto-oncogene in the multiple endocrine neoplasia type 2 syndromes
- 1995 Wessex Regional Genetics Laboratory Seminar, Salisbury, UK: The many faces of *RET*: multiple endocrine neoplasia type 2 and Hirschsprung disease
- 1996 Journées Internationales H P Klotz d'Endocrinologie Clinique, Paris, France: *RET* mutations in multiple endocrine neoplasia type 2 and sporadic medullary thyroid carcinoma
- 1996 Special Seminar, Institut Curie, Paris, France: Mapping of the Cowden disease susceptibility gene: clue to *BRCA3*?
- 1996 Medical Genetics Seminar, Institut Necker, Hopital des Enfants-Malades, Paris, France: Mutations in the *RET* proto-oncogene in MEN 2 and Hirschsprung disease
- 1996 Department of Endocrinology Seminar, King's College Hospital School of Medicine, London, UK: *RET* proto-oncogene in MEN 2 and sporadic MTC
- 1996 Department of Endocrinology Seminar, St. Bartholomew's Hospital, London, UK: Localisation of the gene for Cowden disease: another breast cancer susceptibility gene?
- 1996 Special Seminar, Department of Medical Genetics, Queen's University, Kingston, ON: Cowden syndrome
- 1997 Université Claude Bernard Lyon I, Lyon, France: External examiner, PhD thesis committee (PhD Candidate: Isabelle Schuffenecker)
- 1997 Special Seminar, International Agency for Research on Cancer, Lyon, France: Molecular genetics of Cowden syndrome
- 1997 Special Seminar, Cancer Institute of New Jersey, New Brunswick, NJ: *PTEN* in Cowden syndrome
- 1997 31st Patterson Symposium: Li-Fraumeni syndrome, Manchester, UK: Two-dimensional gene scanning for rapid *p53* mutation detection
- 1997 IV International Thyroid and Neuroendocrine Cancer Workshop, Sicily, Italy: Genotype-phenotype correlations in MEN 2 and genotype-prognosis studies in sporadic medullary thyroid carcinoma
- 1998 Special Seminar, Fox Chase Cancer Center, Philadelphia: *PTEN*, encoding a dual specificity phosphatase, in inherited hamartoma-tumor syndromes
- 1998 Endocrine Grand Rounds, Mt. Sinai Medical Center, NY: The *RET* proto-oncogene in inherited and sporadic medullary thyroid carcinoma
- 1998 Special Seminar, Human Cancer Genetics Program, Comprehensive Cancer Center, Ohio State University, Columbus, OH: The paradox of the *RET* proto-oncogene: multiple endocrine neoplasia and Hirschsprung disease
- 1998 Special Seminar, Human Cancer Genetics Program, MD Anderson Cancer Center, Houston, TX: *PTEN* in inherited hamartoma-tumour syndromes
- 1998 Invited Lecture, First International Lentigenosis Meeting, National Institutes of Health, Bethesda, MD: *PTEN*, Cowden syndrome and Bannayan-Ruvalcaba-Riley syndrome
- 1998 Invited Symposium Lecture, Fourth European Congress of Endocrinology, Seville, Spain: *RET* and *PTEN* mutations in sporadic thyroid tumours
- 1998 Invited Lecture, ASCO Continuing Medical Education Course "Cancer Genetics in Office Practice," Princeton, NJ: Genetics of colorectal cancer
- 1998 Breast Cancer Research Centre, Vancouver, BC: *PTEN* and its role in breast tumourigenesis in Cowden syndrome



- 1998 Invited Lecture, 54<sup>th</sup> Recent Progress in Hormone Research, Skamania Lodge, Stevenson, WA: *PTEN*, encoding a phosphatase, in hereditary and sporadic nonmedullary thyroid tumors
- 1998 Invited Lecture, Gordon Research Conference DNA Alterations in Transformed Cells: New insights into the molecular genetics of cancer, Colby-Sawyer College, NH: *PTEN* mutations in two inherited hamartoma-cancer syndromes and sporadic tumors
- 1998 Invited Lecture, International Congress on Hereditary Cancer Diseases, Düsseldorf, Germany: Cowden syndrome: update on genetic mechanisms and clinical features
- 1998 Grand Rounds, University of Michigan Cancer Center, Ann Arbor, MI: The yin and yang of inherited thyroid cancer
- 1998 Invited Lecture, American Psychological Association Conference on Behavioral Science and Genetics, Tyson's Corner, VA: Genetic testing: from technology to treatment
- 1998 Karolinska Institute, Stockholm, Sweden: Faculty Opponent for PhD Thesis Defence (PhD Candidate: Filip Farnebo)
- 1999 Grand Rounds, NIDDK, NIH, Bethesda, MD: Genetic and epigenetic *PTEN* alterations in inherited and sporadic neoplasia
- 1999 Invited Lectures, NIH-sponsored Phakomatosis Revisited Workshop Rockville, MD: Hamartoses; Cowden syndrome and *PTEN*
- 1999 Invited Lecture, ASCO Train the Trainer Update: Bringing Cancer Genetics to Office Practice, New Orleans, LA: Molecular diagnosis of the inherited hamartoma tumor syndromes
- 1999 Medicine Grand Rounds, Rush Medical School, Chicago, IL: Molecular genetics in office practice: *RET* proto-oncogene mutations in multiple endocrine neoplasia type 2
- 1999 Molecular Medicine Seminar, University of Toronto, Canada: Genetics of *PTEN* in inherited and sporadic cancers
- 1999 Invited Symposium Lecture, American Gastroenterological Association, Orlando, FL: Feast or famine: *RET* proto-oncogene in intestinal ganglioneuromatosis and Hirschsprung disease
- 1999 Invited Plenary Lecture, Seventh International Workshop on Multiple Endocrine Neoplasia, Gubbio, Italy: MEN 2 and the practice of molecular oncology
- 1999 Invited Plenary Lecture, Seventh International Workshop on Multiple Endocrine Neoplasia, Gubbio, Italy: The role of *PTEN* in Cowden syndrome and multiple sporadic cancers

### C. Short Report of Clinical Activities

Description of Clinical Practice: Clinical cancer genetics; medical oncology, especially inherited hamartoma tumor syndromes, and endocrine tumors in a teaching hospital setting.

Patient Load: 20% effort in the practice of clinical cancer genetics. Patients/families seen in cancer genetics clinic are usually complex and labor intensive.

Clinical Contributions: When we and other groups discovered that germline mutations in the *RET* proto-oncogene are associated with MEN 2, clinical diagnostic testing became available within 6 months of our publication. Since then, our work as well as others' work have borne out initial data, such that *RET* testing has now become the clinical standard of care in MEN 2 and all cases of medullary thyroid cancer. Mutation status is important in these entities because it alters clinical

management for the patient and his/her family. I have also worked with at least one CLIA-certified laboratory to ensure quality control and have worked with at least one third party insurer so that *RET* testing is covered 100%.

## Bibliography

### Original Reports:

1. Garber ED, **Eng C**, Puscheck EE, Weil MK, Ward S. Genetics of Ustilago violacea. XII. Half-tetrad analysis and double selection. Bot Gaz 1982; 143:524-9.
2. **Eng CEL**, Strom CM. Analysis of three restriction fragment length polymorphisms in the human type II procollagen gene. Am J Hum Genet 1985; 37:719-732, 1986; 39:122.
3. **Eng CEL**, Strom CM. New syndrome: familial proportionate short stature, intrauterine growth retardation, and recurrent locking of the fingers. Am J Med Genet 1987; 26:217-20.
4. Garber ED, **Eng C**, Stevens DM. Genetics of Ustilago violacea. XXI. Centromere-linkage values and pericentric gene clustering. Curr Genet 1987; 12:555-60.
5. **Eng C**, Aronson MD. Pneumococcal bacteremia in two immunocompetent adults with otitis media and bronchitis. Lancet 1990; 336:1266 / 1991; 337:240-1 (response to critique).
6. **Eng C**, Chopra S. Acute renal failure in nonfulminant hepatitis A. J Clin Gastroenterol 1990; 12:717-8.
7. Banitt P, **Eng C**. Radiculopathy in an elderly woman. Hosp Pract 1991; 26:40.
8. **Eng C**, Korzenik J. Angina pectoris associated with 5-fluorouracil. Hosp Phys 1991; 27:54-7.
9. **Eng C**. Thoracic adenopathy: metastatic seminoma or sarcoid? Hosp Pract 1992; 27:208-10.
10. **Eng C**, Farraye FA, Shulman LN, Peppercorn MA, Krauss CM, Connors JM, Stone RM. The association between myelodysplastic syndromes and Crohn disease. Ann Intern Med 1992; 117:661-2.
11. **Eng C**, Cunningham D, Quade BJ, Schwamm L, Kantoff P, Skarin AT. Meningeal carcinomatosis from transitional cell carcinoma of the bladder. Cancer 1993; 72:553-7.
12. **Eng C**, Li FP, Abramson DH, Ellsworth RM, Wong FL, Goldman MB, Seddon J, Tarbell N, Boice JD, Jr. Mortality from second tumors among long-term survivors of retinoblastoma. J Natl Cancer Inst 1993; 85:1121-8.
13. **Eng C**, Spechler SJ, Ruben R, Li FP. Familial Barrett esophagus and adenocarcinoma of the gastroesophageal junction. Cancer Epidemiol Biomark Prevent 1993; 2:397-9.
14. Mulligan LM, Kwok JBJ, Healey CS, Elsdon MJ, **Eng C**, Gardner E, Love DR, Mole SE, Moore JK, Papi L, Ponder MA, Telenius H, Tunnacliffe A, Ponder BAJ. Germ-line mutations of the *RET* proto-oncogene in multiple endocrine neoplasia type 2A. Nature 1993; 363:458-60.
15. Li FP, **Eng C**. The familial Muir-Torre syndrome. Ann Intern Med 1993; 119:539.

16. Mulligan LM, **Eng C**, Healey CS, Clayton D, Kwok JBJ, Gardner E, Ponder MA, Frilling A, Jackson CE, Lehnert H, Neumann HPH, Thibodeau SN, Ponder BAJ. Specific mutations of the *RET* proto-oncogene are related to disease phenotype in MEN 2A and FMTC. Nature Genet 1994; 6:70-4.
17. **Eng C**, Smith DP, Mulligan LM, Nagai MA, Healey CS, Ponder MA, Gardner E, Scheumann GFW, Jackson CE, Tunnacliffe A, Ponder BAJ. Point mutation within the tyrosine kinase domain of the *RET* proto-oncogene in multiple endocrine neoplasia type 2B and related sporadic tumours. Hum Mol Genet 1994; 3:237-41.
18. **Eng C**, Murday V, Seal S, Mohammed S, Hodgson SV, Chaudary MA, Fentiman I, Ponder BAJ, Eeles RA. Cowden syndrome and Lhermitte-Duclos disease in a family: a single genetic syndrome with pleiotropy? J Med Genet 1994; 31:458-61.
19. Mulligan LM, **Eng C**, Healey CS, Ponder MA, Feldman GL, Li P, Jackson CE, Ponder BAJ. A *de novo* mutation of the *RET* proto-oncogene in a patient with MEN 2A. Hum Mol Genet 1994; 3:1007-8.
20. Attié T, Pelet A, Sarda P, **Eng C**, Edery P, Mulligan LM, Ponder BAJ, Munnich A, Lyonnet S. A 7 bp deletion of the *RET* proto-oncogene in familial Hirschsprung's disease. Hum Mol Genet 1994; 3:1439-40.
21. Gardner E, Mulligan LM, **Eng C**, Healey CS, Kwok JBJ, Ponder MA, Ponder BAJ. Haplotype analysis of MEN 2 mutations. Hum Mol Genet 1994; 3:1771-4.
22. Edery P, Attié T, Mulligan LM, Pelet A, **Eng C**, Ponder BAJ, Munnich A, Lyonnet S. A novel polymorphism in the coding sequence of the human *RET* proto-oncogene. Hum Genet 1994; 94:579-80.
23. Mulligan LM, **Eng C**, Attié T, Lyonnet S, Marsh D, Hyland VJ, Robinson BG, Frilling A, Verellen-Dumoulin C, Safar A, Venter DJ, Munnich A, Ponder BAJ. Diverse phenotypes associated with exon 10 mutations in the *RET* proto-oncogene. Hum Mol Genet 1994; 3:2163-7.
24. Pelet A, Attié T, Goulet O, **Eng C**, Ponder BAJ, Munnich A, Lyonnet S. *De novo* mutations of the *RET* proto-oncogene in Hirschsprung's disease. Lancet 1994; 344:1769-70.
25. **Eng C**, Mulligan LM, Smith DP, Healey CS, Frilling A, Raue F, Neumann HPH, Pfragner R, Behmel A, Lorenzo MJ, Stonehouse TJ, Ponder MA, Ponder BAJ. Mutation of the *RET* proto-oncogene in sporadic medullary thyroid carcinomas. Genes Chrom Cancer 1995; 12:209-12.
26. Songyang Z, Carraway KL III, Eck MJ, Harrison SC, Feldman RA, Mohammadi M, Schlessinger J, Hubbard SR, Smith DP, **Eng C**, Lorenzo MJ, Ponder BAJ, Mayer BJ, Cantley LC. Catalytic site specificity of protein-tyrosine kinases is critical for selective signalling. Nature 1995; 373:536-9.
27. **Eng C**, Smith DP, Mulligan LM, Healey CS, Zvelebil MJ, Stonehouse TJ, Ponder MA, Jackson CE, Waterfield MD, Ponder BAJ. A novel point mutation in the tyrosine kinase domain of the *RET* proto-oncogene in sporadic medullary thyroid carcinoma and in a family with FMTC. Oncogene 1995; 10:509-13.

28. Lorenzo MJ, **Eng C**, Mulligan LM, Stonehouse TJ, Healey CS, Ponder BAJ, Smith DP. Multiple mRNA isoforms of the human *RET* proto-oncogene generated by alternate splicing. Oncogene 1995; 10:1377-83.
29. Frilling A, Höppner W, **Eng C**, Mulligan L, Raue F, Broelsch CE. Presymptomatic genetic screening in families with multiple endocrine neoplasia type 2. J Mol Med 1995; 73:229-33.
30. Frilling A, Höppner W, Raue F, **Eng C**, Mulligan L, Ponder BAJ, Broelsch CE. Spezifische *RET* proto-onkogen mutationen bei verschiedenen hereditären formen des C-zell-karzinoms. Langenbeck Arch Chirurg Forum '95 1995; 303-6.
31. **Eng C**, Mulligan LM, Smith DP, Healey CS, Frilling A, Raue F, Neumann HPH, Ponder MA, Ponder BAJ. Low frequency of germline mutations in the *RET* proto-oncogene in patients with apparently sporadic medullary thyroid carcinoma. Clin Endocrinol 1995; 43:123-7.
32. Attié T, Pelet A, Edery P, **Eng C**, Mulligan LM, Amiel J, Boutrand L, Beldjord C, Nihoul-Fékété C, Munnich A, Ponder BAJ, Lyonnet S. Diversity of *RET* proto-oncogene mutations in familial and sporadic Hirschsprung disease. Hum Mol Genet 1995; 4:1381-6.
33. Crossey PA\*, **Eng C\***, Ginalska-Malinowska M, Lennard TWJ, Wheeler D, Ponder BAJ, Maher ER. Molecular genetic diagnosis of von Hippel-Lindau disease in familial pheochromocytoma. J Med Genet 1995; 32:885-6.
34. Neumann HPH, **Eng C**, Mulligan LM, Glavac D, Zaüner I, Ponder BAJ, Crossey PA, Maher ER, Brauch H. Consequences of direct genetic testing for germline mutations in the clinical management of families with multiple endocrine neoplasia type 2. JAMA 1995; 274:1149-51.
35. Mulligan LM, Marsh DJ, Robinson BG, Schuffenecker I, Zedenius J, Lips CJM, Gagel RF, Takai S-I, Noll WW, Fink M, Raue F, Lacroix A, Thibodeau SN, Frilling A, Ponder BAJ, **Eng C**. Genotype-phenotype correlation in MEN 2: Report of the International *RET* Mutation Consortium. J Intern Med 1995; 238:343-6.
36. Myers SM, **Eng C**, Ponder BAJ, Mulligan LM. Characterization of *RET* proto-oncogene 3' splicing variants and polyadenylation sites: a novel C-terminal for RET. Oncogene 1995; 11:2039-2045.
37. Frilling A, Dralle H, **Eng C**, Raue F, Broelsch CE. Presymptomatic DNA screening in families with multiple endocrine neoplasia type 2 and familial medullary thyroid carcinoma. Surgery 1995; 118:1099-104.
38. Borrello MG, Smith DP, Pasini B, Bongarzone I, Greco A, Lorenzo MJ, Arighi E, Miranda C, **Eng C**, Alberti L, Bocciardi R, Mondellini P, Scopsi L, Romeo G, Ponder BAJ, Pierotti MA. *RET* activation by germline *MEN2A* and *MEN2B* mutations. Oncogene 1995; 11:2419-27.
39. Toogood AA, **Eng C**, Smith DP, Ponder BAJ, Shalet SM. No mutation at codon 918 of the *RET* gene in a family with multiple endocrine neoplasia type 2B. Clin Endocrinol 1995; 43:759-62.

40. **Eng C\***, Crossey PA\*, Mulligan LM, Healey CS, Houghton C, Prowse A, Chew SL, Dahia PLM, O'Riordan JLH, Toledo SPA, Smith DP, Maher ER, Ponder BAJ. Mutations of the *RET* proto-oncogene and the von Hippel-Lindau disease tumour suppressor gene in sporadic and syndromic pheochromocytomas. J Med Genet 1995; 32:934-7.
41. Marsh DJ, Learoyd DL, Andrew SD, Krishnan L, Pojer R, Richardson A-L, Delbridge L, **Eng C**, Robinson BG. Somatic mutations in the *RET* proto-oncogene in sporadic medullary thyroid carcinoma. Clin Endocrinol 1996; 44:249-57.
42. Marsh DJ, Andrew SD, **Eng C**, Learoyd DL, Capes AG, Pojer R, Richardson A-L, Houghton C, Mulligan LM, Ponder BAJ, Robinson BG. Germline and somatic mutations in an oncogene: *RET* mutations in inherited medullary thyroid carcinoma. Cancer Res 1996; 56:1241-3.
43. Neumann HPH, **Eng C**, Mulligan LM. Von Hippel-Lindau disease and pheochromocytoma. JAMA 1996; 275:839-40.
44. **Eng C**, Foster KA, Healey CS, Houghton C, Gayther SA, Mulligan LM, Ponder BAJ. Mutation analysis of the *c-mos* proto-oncogene and the endothelin-B receptor gene in medullary thyroid carcinoma and pheochromocytoma. Br J Cancer 1996; 74:339-41.
45. Edery P, Attié T, Amiel J, Pelet A, **Eng C**, Hofstra RMW, Martelli H, Bidaud C, Munnich A, Lyonnet S. Mutation of the endothelin-3 gene in the Waardenburg-Hirschsprung disease (Shah-Waardenburg syndrome). Nature Genet 1996; 12:442-4.
46. **Eng C**, Mulligan LM, Healey CS, Houghton C, Frilling A, Raue F, Thomas GA, Ponder BAJ. Heterogeneous mutation of the *RET* proto-oncogene in subpopulations of medullary thyroid carcinoma. Cancer Res 1996; 56:2167-70.
47. Nelen MR, Padberg GW, Peeters EAJ, Lin A, van den Helm B, Frants RR, Coulon V, Goldstein AM, van Reen MMM, Easton DF, Eeles RA, Hodgson S, Mulvihill JJ, Murday VA, Tucker MA, Mariman ECM, Starink TM, Ponder BAJ, Ropers HH, Kremer H, Longy M, **Eng C**. Localization of the gene for Cowden disease to 10q22-23. Nature Genet 1996; 13:114-6.
48. van Orsouw NJ, Li D, van der Vlies P, Scheffer H, **Eng C**, Buys CHCM, Li FP, Vijg J. Mutational scanning of large genes by extensive PCR multiplexing and two-dimensional electrophoresis: application to the *RBI* gene. Hum Mol Genet 1996; 5:755-61.
49. Neumann HPH, Bender B, Zäuner I, Berger DP, **Eng C**, Brauch H, Zbar B. Monogenetic hypertension and pheochromocytoma. Am J Kidney Dis 1996; 28:329-33.
50. Salomon R, Attié T, Pelet A, Bidaud C, **Eng C**, Amiel J, Sarnacki S, Goulet O, Ricour C, Nihoul-Fékété C, Munnich A, Lyonnet S. Germline mutations of a RET ligand, glial cell line-derived neurotrophic factor, are not sufficient to cause Hirschsprung disease. Nature Genet 1996; 14:345-347.
51. **Eng C**, Clayton D, Schuffenecker I, Lenoir G, Cote G, Gagel RF, Ploos van Amstel HK, Lips CJM, Nishisho I, Takai S-I, Marsh DJ, Robinson BG, Frank-Raue K, Raue F, Xue F, Noll WW, Romei C, Pacini F, Fink M, Niederle B, Zedenius J, Nordenskjöld M, Komminoth P, Hendy GN, Gharib H, Thibodeau SN, Lacroix A, Frilling A, Ponder BAJ, Mulligan LM. The relationship between specific *RET* proto-oncogene mutations and

- disease phenotype in multiple endocrine neoplasia type 2. International *RET* Mutation Consortium analysis. *JAMA* 1996; 276:1575-9.
52. Ivanchuk SM, Myers SM, **Eng C**, Mulligan LM. *De novo* mutation of GDNF, ligand for the RET/GDNFR- $\alpha$  receptor complex, in Hirschsprung disease. *Hum Mol Genet* 1996; 5:2023-6.
  53. Schuffenecker I, Ginet N, Goldgar D, **Eng C**, Chambe B, Boneu A, Houdent C, Pallo D, Schlumberger M, Thivolet C, Lenoir GM. Prevalence and parental origin of *de novo RET* mutations in multiple endocrine neoplasia type 2A and familial medullary thyroid carcinoma. *Am J Hum Genet* 1997; 60:233-7.
  54. Kerangueven F, Eisinger F, Noguchi T, Allione F, Wargniez V, **Eng C**, Padberg G, Theillet C, Jacquemier J, Longy M, Sobol H, Birnbaum D. Loss of heterozygosity in human breast carcinomas in the ataxia telangiectasia, Cowden's disease and *BRCA1* gene regions. *Oncogene* 1997; 14:339-47.
  55. Dahia PLM, Toledo SPA, Mulligan LM, Maher ER, Grossman AB, **Eng C**. Mutation analysis of glial cell line-derived neurotrophic factor (GDNF), a ligand for the RET/GDNF receptor  $\alpha$  complex, in sporadic pheochromocytomas. *Cancer Res* 1997; 57:310-3.
  56. Boccia LM, Green JS, Joyce C, **Eng C**, Taylor SAM, Mulligan LM. Mutations of *RET* codon 768 is associated with the FMTC phenotype. *Clin Genet* 1997; 51:81-5.
  57. Marsh DJ, Zheng Z, Zedenius J, Kremer H, Padberg GW, Larsson C, Longy M, **Eng C**. Differential loss of heterozygosity in the region of the Cowden locus within 10q22-23 in follicular thyroid adenomas and carcinomas. *Cancer Res* 1997; 57:500-3.
  58. Marsh DJ, Andrew SD, Learoyd DL, Pojer R, **Eng C**, Robinson BG. Deletion-insertion mutation encompassing *RET* codon 634 is associated with medullary thyroid carcinoma. *Hum Mutat* 1997; Mutations in Brief #12 Online.
  59. Ivanchuk SM, **Eng C**, Cavenee WK, Mulligan LM. The expression of *RET* and its multiple splice forms in developing human kidney. *Oncogene* 1997; 14:1811-8.
  60. Liaw D, Marsh DJ, Li J, Dahia PLM, Wang SI, Zheng Z, Bose S, Call KM, Tsou HC, Peacocke M, **Eng C\***, Parsons R\*. Germline mutations of the *PTEN* gene in Cowden disease, an inherited breast and thyroid cancer syndrome. *Nature Genet* 1997; 16:64-7. (\*Joint Senior Authors)
  61. Woodward ER, **Eng C**, McMahon R, Voutilainen R, Affara NA, Ponder BAJ, Maher ER. Genetic predisposition to pheochromocytoma: analysis of candidate genes *GDNF*, *RET* and *VHL*. *Hum Mol Genet* 1997; 6:1051-6.
  62. Bidaud C, Salomon R, Vancamp G, Pelet A, Attié T, **Eng C**, Bonduelle M, Amiel J, Nihoulfékéte C, Willems PJ, Munnich A, Lyonnet. Endothelin-3 gene mutations in isolated and syndromic Hirschsprung disease. *Eur J Hum Genet* 1997; 5:247-51.
  63. Marsh DJ, Dahia PLM, Zheng Z, Liaw D, Parsons R, Gorlin RJ, **Eng C**. Germline mutations in *PTEN* are present in Bannayan-Zonana syndrome. *Nature Genet* 1997; 16:333-4.

64. Myers MP, Stolarov J, **Eng C**, Li J, Wang SI, Wigler MH, Parsons R, Tonks NK. PTEN, the tumor suppressor from human chromosome 10q23, is a dual specificity phosphatase. Proc Natl Acad Sci (USA) 1997; 94:9052-7.
65. Marsh DJ, Zheng Z, Arnold A, Andrew SD, Learoyd D, Frilling A, Komminoth P, Neumann HPH, Ponder BAJ, Rollins BJ, Shapiro GI, Robinson BG, Mulligan LM, **Eng C**. Mutation analysis of glial cell line-derived neurotrophic factor (*GDNF*), a ligand for the RET/co-receptor complex, in MEN 2 and sporadic neuroendocrine tumors. J Clin Endocrinol Metab 1997; 82:3025-8.
66. Tsou HC, Teng D, Ping XL, Broncolini V, Davis T, Hu R, Xie XX, Gruener AC, Schrager CA, Christiano AM, **Eng C**, Steck P, Ott J, Tavtigian S, Peacocke M. Role of *MMAC1* mutations in early onset breast cancer: causative in association with Cowden syndrome and excluded in *BRCA1*-negative cases. Am J Hum Genet 1997; 61:1036-43.
67. Gimm O, Marsh DJ, Andrew SD, Frilling A, Dahia PLM, Mulligan LM, Zajac JD, Robinson BG, **Eng C**. Germline dinucleotide mutation in codon 883 of the *RET* proto-oncogene in multiple endocrine neoplasia type 2B without codon 918 mutation. J Clin Endocrinol Metab 1997; 82:3902-4.
68. Dahia PLM, Marsh DJ, Zheng Z, Zedenius J, Komminoth P, Frisk T, Wallin G, Parsons R, Longy M, Larsson C, **Eng C**. Somatic deletions and mutations of *PTEN* in sporadic thyroid tumors. Cancer Res 1997; 57:4710-3.
69. Marsh DJ, Roth S, Lunetta K, Hemminki A, Dahia PLM, Sistonen P, Zheng Z, Caron S, van Orsouw NJ, Bodmer WF, Cottrell S, Dunlop MG, Eccles D, Hodgson SV, Järvinen H, Kellokumpu I, Markie D, Neale K, Phillips R, Rozen P, Syngal S, Vijn J, Tomlinson IPM, Aaltonen LA, **Eng C**. Exclusion of *PTEN* and 10q22-24 as the susceptibility locus for juvenile polyposis syndrome (JPS). Cancer Res 1997; 57:5017-21.
70. Marsh DJ, Dahia PLM, Coulon V, Zheng Z, Dorion-Bonnet F, Call KM, Little R, Lin AY, Goldstein A, Eeles RA, Hodgson SV, Richardson A-L, Robinson BG, Weber HC, Longy M, **Eng C**. Allelic imbalance, including deletion of *PTEN/MMAC1*, at the Cowden disease locus on 10q22-23 in hamartomas from patients with Cowden disease and germline *PTEN* mutation. Genes Chrom Cancer 1998; 21:61-9.
71. Gordon CM, Majzoub JA, Marsh DJ, Mulliken J, Ponder BAJ, Robinson BG, **Eng C**. Four cases of mucosal neuroma syndrome: MEN 2B or not 2B? J Clin Endocrinol Metab 1998; 83:17-20.
72. Schuffenecker I, Virally-Monod M, Brohet R, Goldgar D, Conte-Devolx B, Leclerc L, Chabre O, Boneu A, Caron J, Houdent C, Modigliani E, Rohmer V, Schlumberger M, **Eng C**, Guillausseau PJ, Lenoir G. Risk and penetrance of primary hyperparathyroidism in MEN 2A families with codon 634 mutations of the *RET* proto-oncogene. J Clin Endocrinol Metab 1998; 83:487-91.
73. **Eng C**, Myers SM, Kogon MD, Sanicola M, Hession C, Cate RL, Mulligan LM. Genomic structure and chromosomal localisation of the human *GDNFR- $\alpha$*  gene. Oncogene 1998; 16:597-601.



74. Peters N, Wellenreuther R, Rollbrocker B, Hayashi Y, Meyer-Puttlitz B, Dürr E-M, Lenartz D, Marsh DJ, Schramm J, Wiestler OD, Parsons R, **Eng C**, von Deimling A. Analysis of the *PTEN* gene in human meningiomas. Neuropathol Appl Neurobiol 1998; 24:3-8.
75. Marsh DJ, Coulon V, Lunetta KL, Rocca-Serra P, Dahia PLM, Zheng Z, Liaw D, Caron S, Duboué B, Lin AY, Richardson AL, Bonnetblanc J-M, Bressieux J-M, Cabarrot-Moreau A, Chompret A, Demange L, Eeles RA, Yahanda AM, Fearon ER, Fricker JP, Gorlin RJ, Hodgson SV, Huson S, Lacombe D, LePrat F, Odent S, Toulouse C, Olapade OI, Sobol H, Tishler S, Woods CG, Robinson BG, Weber HC, Parsons R, Peacocke M, Longy M, **Eng C**. Mutation spectrum and genotype-phenotype analyses in Cowden disease and Bannayan-Zonana syndrome, two hamartoma syndromes with germline *PTEN* mutation. Hum Mol Genet 1998; 7:507-15.
76. Feilotter HE, Nagai MA, Boag AH, **Eng C**, Mulligan LM. Analysis of *PTEN* and the 10q23 region in primary prostate carcinomas. Oncogene 1998; 16:1743-8.
77. Mulligan LM, Timmer T, Ivanchuk SM, Campling BG, Sundaresan V, Rabbitts PH, Hofstra RMW, **Eng C**. Investigations of the genes encoding RET and its ligand complex, GDNF/GFR $\alpha$ -1, in small cell lung carcinoma. Genes Chrom Cancer 1998; 21:326-32.
78. Dürr E-M, Rollbrocker B, Hayashi Y, Peters N, Meyer-Puttlitz B, Louis DN, Schramm J, Wiestler OD, Parsons R, **Eng C**, von Deimling A. *PTEN* mutations in gliomas and glioneuronal tumors. Oncogene 1998; 16:2259-2264.
79. van Orsouw NJ, Dhanda RK, Rines RD, Smith WM, Sigalas I, **Eng C**, Vijg J. Rapid design of denaturing gradient-based two-dimensional electrophoretic gene mutation scanning tests. Nucleic Acid Res 1998; 26:2398-2406.
80. Dahia PLM, FitzGerald MG, Zhang X, Marsh DJ, Zheng Z, Pietsch T, von Deimling A, Haluska FG, Haber DA, **Eng C**. A highly conserved processed *PTEN* pseudogene is located on chromosome band 9p21. Oncogene 1998; 16:2403-6.
81. Pesche S, Latil A, Muzeau F, Cussenot O, Fournier G, Longy M, **Eng C**, Lidereau R. *PTEN/MMAC1/TEP1* involvement in primary prostate cancers. Oncogene 1998; 16:2879-83.
82. Rines RD, van Orsouw NJ, Sigalas I, Li FP, **Eng C**, Vijg J. Comprehensive mutational analysis of *p53* coding region by two-dimensional gene scanning. Carcinogenesis 1998; 19:979-84.
83. **Eng C**, Peacocke M. *PTEN* mutation analysis as a molecular diagnostic tool in the inherited hamartoma-cancer syndromes. Nature Genet 19:223.
84. Svensson P-J, Molander M-L, **Eng C**, Anvret M, Nordenskjöld A. Low frequency of *RET* mutations in Hirschsprung disease in Sweden. Clin Genet 1998; 54:39-44.
85. Stratakis CA, Kirschner LS, Taymans SE, Tomlinson IPM, Marsh DJ, Torpy DJ, Eccles DM, Theaker J, Houlston RS, Blouin J-L, Antonarakis SE, Basson CT, **Eng C**, Carney JA. Carney complex, Peutz-Jeghers syndrome, Cowden disease, and Bannayan-Zonana syndrome share cutaneous and endocrine manifestations but not genetic loci. J Clin Endocrinol Metab 1998; 83:2972-6.

86. FitzGerald MG, Marsh DJ, Wahrer D, Caron S, Bell S, Shannon KEM, Ishioka C, Isselbacher KJ, Garber JE, **Eng C**, Haber DA. Germline mutations in *PTEN* are an infrequent cause of genetic predisposition to breast cancer. Oncogene 1998; 17:727-32.
87. Borrego S, **Eng C**, Sánchez B, Sáez M-E, Navarro E, Antinolo G. Molecular analysis of the *RET* and *GDNF* genes in a family with multiple endocrine neoplasia type 2A and Hirschsprung disease. J Clin Endocrinol Metab 1998; 83:3361-4.
88. Dhanda RK, van Orsouw NJ, Sigalas I, **Eng C**, Vijg J. Critical factors in the performance and cost of two-dimensional gene scanning: *RBI* as a model. BioTechniques 1998; 25:664-75.
89. Somerville RPT, Shoshan Y, **Eng C**, Barnett G, Miller D, Cowell JK. Molecular analysis of two putative tumour suppressor genes, *PTEN* and *DMBT*, which have been implicated in glioblastoma multiforme disease progression. Oncogene 1998; 17:1755-1757.
90. Longy M, Coulon V, Duboué B, David A, Larrègue M, **Eng C**, Amati P, Kraitsirps J-L, Bottani A, Lacombe D, Bonneau D. Mutations of *PTEN* in patients with Bannayan-Riley-Ruvalcaba phenotype. J Med Genet 1998; 35:886-9.
91. Smith WM, van Orsouw NJ, Fox EA, Kolodner RD, Vijg J, **Eng C**. Accurate, high throughput "snapshot" detection of *hMLH1* mutations by two-dimensional electrophoresis. Genet Testing 1998; 2:43-53.
92. Dhanda RK, Smith WM, Scott CB, **Eng C**, Vijg J. Automated two-dimensional DNA electrophoresis: application to genetic testing. Genet Testing 1998; 2:67-70.
93. Dabora SL, Sigalas I, Hall F, **Eng C**, Vijg J, Kwiatkowski DJ. Comprehensive mutation analysis of *TSC1* with two-dimensional DNA electrophoresis with DGGE. Ann Hum Genet 1998; 62:491-504.
94. Marsh DJ, Dahia PLM, Caron S, Kum JB, Frayling IM, Tomlinson IPM, Hughes KS, Hodgson SV, Murday VA, Houlston R, **Eng C**. Germline *PTEN* mutations in Cowden syndrome-like families. J Med Genet 1998; 35:881-5.
95. Houlston R, Bevan S, Williams A, Young J, Dunlop M, Rozen P, **Eng C**, Markie D, Woodford-Richens K, Rodriguez-Bigas M, Leggett B, Neale K, Phillips R, Sheridan E, Hodgson S, Twama T, Eccles D, Fagan K, Bodmer W, Tomlinson I. Mutations in *DPC4* (*SMAD4*) cause juvenile polyposis syndrome, but only account for a minority of cases. Hum Mol Genet 1998; 7:1907-1912.
96. **Eng C**, Thomas GA, Neuberger DS, Mulligan LM, Healey CS, Houghton C, Frilling A, Raue F, Williams ED, Ponder BAJ. Mutation of the *RET* proto-oncogene is correlated with RET immunostaining in subpopulations of cells in sporadic medullary thyroid carcinoma. J Clin Endocrinol Metab 1998; 83:4310-3.
97. **Eng C**, Marsh DJ, Robinson BG, Chow CW, Patton MA, Southey MC, Venter DJ, Ponder BAJ, Milla PJ, Smith VV. Germline *RET* codon 918 mutation in apparently isolated intestinal ganglioneuromatosis. J Clin Endocrinol Metab 1998; 83:4191-4.
98. Zori RT, Marsh DJ, Graham GE, Marliiss EB, **Eng C**. Germline *PTEN* mutation in a family with Cowden syndrome and Bannayan-Ruvalcaba-Riley syndrome. Am J Med Genet 1998; 80:399-402.

99. Feilotter HE, Coulon V, McVeigh JL, Boag AH, Dorion-Bonnet F, Duboué B, Latham WCW, **Eng C**, Mulligan LM, Longy M. Analysis of the 10q23 chromosomal region and the *PTEN* gene in human sporadic breast carcinoma. Br J Cancer 1999; 79:718-23.
100. Myers SM, Salomon R, Gössling A, Pelet A, **Eng C**, von Deimling A, Lyonnet S, Mulligan LM. Absence of germline *GFR $\alpha$ -1* mutations in Hirschsprung disease. J Med Genet 1999; 36:217-20.
101. Dahia PLM, Aguiar RCT, Alberta J, Kum JB, Caron S, Sills H, Marsh DJ, Freedman A, Ritz J, Stiles C, **Eng C**. PTEN is inversely correlated with the cell survival factor PKB/Akt and is inactivated by diverse mechanisms in haematologic malignancies. Hum Mol Genet 1999; 8:185-93.
102. Svensson PJ, Tapper-Persson M, Anvret M, Molander M-L, **Eng C**, Nordenskjöld A. Mutations in the endothelin-receptor  $\beta$  gene in Hirschsprung disease in Sweden. Clin Genet 1999; 55:215-7.
103. Gimm O, Neuberg DS, Marsh, DJ, Dahia PLM, Hoang-Vu C, Raue F, Hinze R, Dralle H, **Eng C**. Over-representation of a germline *RET* sequence variant in patients with sporadic medullary thyroid carcinoma and somatic *RET* codon 918 mutation. Oncogene 1999; 18:1369-74.
104. Otto LR, Boriack RL, Marsh DJ, Kum JB, **Eng C**, Burlina AB, Bennett MJ. Long-chain L-3-hydroxyacyl-coA dehydrogenase (LCHAD) deficiency does not appear to be the primary cause of lipid myopathy in patients with Bannayan-Riley-Ruvalcaba syndrome. Am J Med Genet 1999; 83:3-5.
105. Gimm O, Gössling A, Marsh DJ, Dahia PLM, Myers SM, Mulligan LM, von Deimling A, **Eng C**. Somatic deletion of the gene encoding GFR $\alpha$ -1, a co-receptor of RET, in sporadic brain tumours. Br J Cancer 1999; 80:383-6.
106. Nilsson O, Tisell L-E, Jansson S, Ahlman H, Gimm O, **Eng C**. Adrenal and extra-adrenal pheochromocytomas in a family with germline *RET* V804L mutation. JAMA 1999; 281:1587-8.
107. Vestergaard P, Kroustrup JP, Ronne H, **Eng C**, Laurberg P. Neuromas in multiple endocrine neoplasia type 2A with a *RET* codon 611 mutation. J Endo Genet 1999; 1:33-8.
108. Shannon KE, Gimm O, Hinze R, Dralle H, **Eng C**. Germline V804M mutation in the *RET* proto-oncogene in two apparently sporadic cases of MTC presenting in the seventh decade of life. J Endo Genet 1999; 1:39-46.
109. Laugé A, Lefèbvre C, Laurent-Puig P, Gad S, **Eng C**, Longy M, Stoppa-Lyonnet D. No evidence for germline *PTEN* mutations in families with breast and brain tumours. Intl J Cancer 1999; 84:216-9.
110. Smith VV, **Eng C**, Milla PJ. Intestinal ganglioneuromatosis and multiple endocrine neoplasia type 2B: implications for treatment. Gut 1999; 45:143-6.

111. Sarraf P, Mueller E, Smith WM, Wright HM, Kum JB, Aaltonen LA, de la Chapelle A, Speigelman BM, **Eng C**. Loss-of-function mutations in *PPAR $\gamma$*  associated with human colon cancer. Mol Cell 1999; 3:799-804.
112. Syngal S, Fox EA, Davidio M, Li C, **Eng C**, Kolodner RD, Garber JE. Interpretation of genetic test results for hereditary nonpolyposis colorectal cancer. Implications for predisposition testing. JAMA 1999; 282:247-253.
113. Bevan S, Woodford-Richens K, Rozen P, **Eng C**, Young J, Dunlop M, Neale K, Phillips R, Markie D, Rodriguez-Bigas M, Leggett B, Sheridan E, Hodgson S, Iwama T, Eccles D, Bodmer W, Houlston R, Tomlinson I. Screening *SMAD1*, *SMAD2*, *SMAD3* and *SMAD5* for germline mutations in juvenile polyposis syndrome. Gut 1999; 45:406-8.
114. Clifford SC, Walsh S, Hewson K, Green EK, Brinke A, Green PM, Gianelli F, **Eng C**, Maher ER. Genomic organisation and chromosomal localisation of the human *CUL2* gene and the role of von Hippel-Lindau tumor suppressor-binding protein (CUL2 and VBP-1) mutation and loss in renal cell carcinoma development. Gene Chrom Cancer 1999; 26:20-8.
115. Gimm O, Greco A, Hinze R, Dralle H, Pierotti M, **Eng C**. Sequence variants in *NTRK1* in human sporadic medullary thyroid carcinomas. J Clin Endocrinol Metab 1999; 84:2784-7.
116. Marsh DJ, Kum JB, Lunetta KL, Bennett MJ, Gorlin RJ, Ahmed SF, Bodurtha J, Crowe C, Curtis MA, Dasouki M, Dunn T, Feit H, Geraghty MT, Graham JM, Hodgson SV, Hunter A, Korf BR, Manchester D, Miesfeldt S, Murday VA, Nathanson KA, Parisi M, Pober B, Romano C, Tolmie JL, Trembath R, Winter RM, Zackai EH, Zori RT, Weng LP, Dahia PLM, **Eng C**. *PTEN* mutation spectrum and genotype-phenotype correlations in Bannayan-Riley-Ruvalcaba syndrome suggest a single entity with Cowden syndrome. Hum Mol Genet 1999; 8:1461-72.
117. Duerr EM, Gimm O, Kum JB, Clifford SC, Toledo SPA, Maher ER, Dahia PLM, **Eng C**. Differences in allelic distribution of two polymorphisms in the VHL-associated gene *CUL2* in pheochromocytoma patients without somatic *CUL2* mutations. J Clin Endocrinol Metab 1999; 84:3207-11.
118. van Orsouw NJ, Dhanda RK, Elhaji Y, Narod SA, Li FP, **Eng C**, Vijg J. A highly accurate, low cost test for *BRCA1* mutations. J Med Genet 1999; 36:747-753.
119. Borrego S, Saez ME, Ruiz A, Gimm O, López-Alonzo M, Antiñolo G, **Eng C**. Specific sequence polymorphisms in the *RET* proto-oncogene are over-represented in individuals with Hirschsprung disease and may represent loci modifying phenotypic expression. J Med Genet 1999; 36:771-774.
120. Perren A, Weng LP, Boag AH, Ziebold U, Kum JB, Dahia PLM, Komminoth P, Lees JA, Mulligan LM, Mutter GL, **Eng C**. Immunocytochemical evidence of loss of PTEN expression in primary ductal adenocarcinomas of the breast. Am J Pathol (in press)
121. Ahmed SF, Marsh DJ, Weremowicz S, Morton CC, Williams DM, **Eng C**. Balanced translocation of 10q and 13q, including the *PTEN* gene, in a boy with an HCG-secreting tumor and the Bannayan-Riley-Ruvalcaba syndrome. J Clin Endocrinol Metab (in press)
122. Yeh JJ, Marsh DJ, Zedenius J, Dwight T, Delbridge L, Robinson BG, **Eng C**. Fine structure deletion analysis of 10q22-24 demonstrates novel regions of loss and suggests

that sporadic follicular thyroid adenomas and follicular thyroid carcinomas develop along distinct parallel neoplastic pathways. Gene Chrom Cancer (in press)

123. Weng LP, Smith WM, Dahia PLM, Ziebold U, Gil E, Lees JA, **Eng C**. PTEN suppresses breast cancer cell growth by phosphatase activity-dependent G1 arrest followed by cell death. Cancer Res (in press)

## Reviews:

1. **Eng C**, Blackstone MO. The Peutz-Jeghers syndrome. Med Rounds 1988; 1:165-71.
2. **Eng C**, Skolnick AE, Come SE. Elevated creatine kinase and malignancy. Hosp Pract 1990; 25:123-30.
3. **Eng C**, Ponder BAJ. The role of gene mutations in the genesis of familial cancers. FASEB J 1993; 7:910-9.
4. **Eng C**, Stratton M, Ponder B, Murday V, Easton D, Sacks N, Watson M, Eeles R. Familial cancer syndromes. Lancet 1994; 343:709-13.
5. Smith DP, **Eng C**, Ponder BAJ. Germline mutations of the *RET* proto-oncogene in the multiple endocrine neoplasia type 2 syndromes and Hirschsprung disease. J Cell Science 1994; suppl 18:43-9.
6. Chew SL, **Eng C**. Multiple endocrine neoplasia type 2 and related genetic conditions. Curr Opin Endocrinol Diabet 1995; 2:121-6.
7. Reynolds LF, **Eng C**. *RET* proto-oncogene mutations in multiple endocrine neoplasia type 2 and Hirschsprung disease. Curr Opin Pediatr 1995; 7:702-9.
8. **Eng C**. Seminars in Medicine of the Beth Israel Hospital, Boston: *RET* proto-oncogene mutations in multiple endocrine neoplasia type 2 and Hirschsprung disease. N Engl J Med 1996; 335:943-51.
9. **Eng C**, Mulligan LM. Mutations of the *RET* proto-oncogene in the multiple endocrine neoplasia type 2 syndromes, related sporadic tumours and Hirschsprung disease. Hum Mutat 1997; 9:97-109.
10. **Eng C**. Predictive testing in hereditary medullary thyroid carcinoma. Acta Chir Austriaca 1997; 29:5-8.
11. Marsh DJ, Mulligan LM, **Eng C**. *RET* proto-oncogene mutations in MEN 2 and medullary thyroid carcinoma. Horm Res 1997; 47:168-78.
12. **Eng C**, Vijg J. Genetic testing: the problems and the promise. Nature Biotechnol 1997; 15:422-6.
13. **Eng C**, Schneider KA, Fraumeni JF Jr, Li FP. Meeting report: third international workshop on collaborative multidisciplinary studies of *p53* and other predisposing genes in Li-Fraumeni syndrome. Cancer Epidemiol Biomark Prevent 1997; 6:379-83.
14. Edery P, **Eng C**, Munnich A, Lyonnet S. RET in human development and organogenesis. BioEssays 1997; 19:389-95.
15. Neumann HPH, Bender BU, Januszewicz A, Janetschek G, **Eng C**. Phéochromocytome familial. Act Néphrol 1997; 337-50.
16. **Eng C**. Cowden syndrome. J Genet Counsel 1997; 6:181-91.

17. **Eng C.** From bench to bedside ... but when? Genome Res 1997; 7:669-72.
18. Antonarakis SE, Ashburner M, Auerback AD, Beaudet AL, Beckmann JS, Beutler E, Cooper DN, Cotton RGH, den Dunnen JT, Desnick RJ, **Eng C**, Fasman KH, Goldman D, Hayashi K, Hutchinson F, Kazazian HH, Keen J, King MC, Lehvaslaiho H, McAlpine, PJ, McKusick V, Motulski AG, Povey S, Schorderet DF, Scriver CR, Shows TB, Superti-Furga A, Tay AHN, Tsui L-C, Valle D, Vihinen M. Recommendations for a nomenclature system for human gene mutations. Hum Mutat 1998; 11:1-3.
19. **Eng C.** Genetics of Cowden syndrome: through the looking glass of oncology. Intl J Oncol 1998; 12:701-10.
20. **Eng C.** A novel tumor suppressor gene on chromosome 10 involved in endocrine neoplasia. Curr Opin Endocrinol Diabet 1998; 5:40-8.
21. **Eng C**, Ji H. Invited Editorial. Molecular classification of the inherited hamartoma polyposis syndromes: clearing the muddy waters. Am J Hum Genet 1998; 62:1020-2.
22. **Eng C.** *RET* proto-oncogene in the development of human cancer. J Clin Oncol 1999; 17:380-93.
23. Iliopoulos O, **Eng C.** Genetic and clinical aspects of familial renal neoplasms. Sem Oncol (in press)

#### **Books and Other Monographs:**

1. **Eng C.** The molecular genetics of the human type II procollagen gene and its application to the study of the chondrodystrophies. PhD dissertation, University of Chicago, 1986.
2. Ponder BAJ, **Eng C**, Smith DP. Mutations of the *RET* proto-oncogene in the multiple endocrine neoplasia type 2 syndromes and Hirschsprung disease. Accomplishments in Cancer Research 1994, GM Cancer Research Foundation, New York, 1994:118-32.
3. Neumann HPH, **Eng C**, Bender BU. Multiple endocrine neoplasia type 2. In: Sznjderman M, Januszewicz W, Januszewicz A, ed, Hormonal Hypertension, Wydawnictwo Naukowe, Warsaw, 1997: 292-308.
4. Maher ER, **Eng C.** Genetics of pheochromocytoma. In: Thakker RV, ed, Molecular Genetics of Endocrine Disorders, Chapman and Hall, London, 1997: 279-89.
5. **Eng C**, Ponder BAJ. Multiple endocrine neoplasia type 2 and medullary carcinoma of the thyroid. In: Grossman A, ed, Clinical Endocrinology, 2nd ed, Blackwell Science, Oxford, 1998:635-53.
6. **Eng C**, Parsons R. Cowden syndrome. In: Vogelstein B, Kinzler KW, ed, The Genetic Basis of Human Cancer, McGraw Hill and Co., New York, 1998: 519-25.
7. de la Chapelle A, **Eng C.** Molecular genetic diagnosis in hereditary cancer. In: Perry MC, ed, ASCO Educational Book, Lippincott, Williams and Wilkins, Baltimore, MD, 1999:445-453.

8. **Eng C**, Maher ER. Dominant genes and phacomatoses associated with multiple primary cancers. In: Neugut AI, Robinson E, Meadows AT, eds, Multiple Primary Cancers, Lippincott, Williams and Wilkins, Philadelphia, 1999: 165-196.
9. **Eng C**, Parsons P. Cowden syndrome. In: Scriver CR, Beaudet AL, Sly WS, Valle D, Childs B, Vogelstein B, eds, The Metabolic and Molecular Bases of Inherited Disease, 8th ed, McGraw Hill and Co., New York (in press)
10. Gimm O, **Eng C**. Medullary thyroid carcinoma. In: Souhami RL, Tannock I, Hohenberger P, Horiot J-C, eds, Oxford Textbook of Oncology, 2nd ed, Oxford University Press, Oxford (in press)
11. **Eng C**. Cowden syndrome. In: Thakker RV, Wiersinga W, eds, Oxford Textbook of Endocrinology, Oxford University Press, Oxford (in press)
12. Mulligan LM, **Eng C**. Dissecting the genetics of the *RET* proto-oncogene: paradigm for the practice of molecular medicine. In: Ehrlich M, ed, DNA Alterations in Cancer: Genetic and Epigenetic Changes, BioTechniques Books, Natick, MA (in press)
13. **Eng C**, Burt R, Talbot I. Cowden syndrome. In: Hamilton S, Aaltonen L, eds, Pathology and molecular genetics of digestive tract tumours (WHO Classification of Tumours), IARC Press, Lyon (in press)
14. Dahia PLM, **Eng C**, Eds, Genetic Disorders of Endocrine Neoplasia, Karger, Basel (in preparation)



## Abstracts:

1. **Eng CEL**, Strom CM. Analysis of HindIII polymorphisms in the human type II collagen gene in achondroplasia. Pediatr Res 1985; 19:247A.
2. Strom CM, **Eng CEL**, Christides T, Belles C, Pauli R. Detection of gene deletions in the human type II collagen gene in 8 patients with achondroplasia using gene dosage analysis. Pediatr Res 1985; 19:254A.
3. **Eng CEL**, Strom CM. Difference in type II procollagen genotype distribution between patients with sporadic achondroplasia and their normal parents. Am J Hum Genet 1985; 37:A152.
4. Strom CM, **Eng CEL**. Deletion of the promotor and the majority of the type II procollagen gene in 2 patients with achondroplasia. Am J Hum Genet 1985; 37:A177.
5. Vitek CR, Strom CM, Fee M, **Eng CEL**, Rich SS, Orr HT, Gorlin RJ, Whitley CB. Analysis of type II procollagen gene in skeletal dysplasias and identification of a new, rare RFLP. Am J Hum Genet 1985; 37:A180.
6. **Eng C**. Organization of the mitochondrial genome in the fungal genus Ustilago. Medicine on the Midway 1988; 42:14.
7. **Eng C**, Abramson DH, Ellsworth RM, Boice JD, Jr, Seddon J, Goldman M, Li FP. Causes of late mortality in retinoblastoma patients. Proc Am Assoc Cancer Res 1991; 32:221 (Abs 1317).
8. Ponder BAJ, Mulligan LM, Kwok JB, **Eng C**, Tunnacliffe AG, Ponder MA, Gardner E, Moore JK, Healey K, Elsdon MJ. The molecular genetics of multiple endocrine neoplasia type 2 (MEN 2). Interactions of Cancer Susceptibility Genes and Environmental Carcinogens (AACR/IARC). Lyon, France, 1993.
9. Neumann HPH, Mulligan LM, Kwok JBJ, **Eng C**, Ponder BAJ. Germ-line mutations of the *RET* proto-oncogene in multiple endocrine neoplasia type 2A. Deutsche Gesellschaft Endokrinol, March, 1994.
10. Ponder B, Mulligan L, Kwok J, **Eng C**, Tunnacliffe A, Ponder M, Gardner E, Moore J, Healey K, Elsdon M. The molecular genetics of multiple endocrine neoplasia-2 (MEN 2). J Cell Biochem 1994; S18D:98.
11. **Eng C**, Mulligan LM, Smith DP, Kwok JBJ, Nagai MA, Healey CS, Ponder MA, Gardner E, Moore JK, Elsdon MJ, Tunnacliffe A, Ponder BAJ. The molecular genetics of multiple endocrine neoplasia type 2. March of Dimes 25th Clinical Genetics Conference, Orlando, FL, March, 1994.
12. Ponder B, Mulligan L, **Eng C**, Edery P, Lyonnet S, Smith D, Tunnacliffe A, Kwok J. The multiple endocrine neoplasia type 2 syndromes and Hirschsprung disease are due to different mutations in the receptor tyrosine kinase RET. Fourth European Workshop on Cytogenetics and Molecular Genetics of Human Solid Tumors, Netherlands, April, 1994.

13. Lyonnet S, Attié T, Pelet A, **Eng C**, Edery P, Nihoul-Fékété C, Ponder BAJ, Munnich A. Spectrum of mutations of the *ret* proto-oncogene in Hirschsprung's disease. Am J Hum Genet 1994; 55 (suppl):A6 (Abstract 19).
14. Attié T, **Eng C**, Mulligan LM, Edery P, Verellen C, Ponder BAJ, Munnich A, Lyonnet S. Mutations in exon 10 of the *ret* proto-oncogene in Hirschsprung's disease. Am J Hum Genet 1994; 55 (suppl):A210 (Abstract 1223).
15. **Eng C**, Mulligan LM, Lyonnet S, Edery P, Smith DP, Kwok JBJ, Gardner E, Healey CS, Ponder MA, Munnich A, Ponder BAJ. Mutations of the *RET* proto-oncogene in the multiple endocrine neoplasia type 2 syndromes and Hirschsprung disease. J Med Genet 1995; 32:135.
16. Binchy A, Evans G, **Eng C**, Ponder B, Crauford D. Factors influencing decisions on whether to proceed with predictive testing for breast/ovarian cancers. J Med Genet 1995; 32:140.
17. **Eng C**, Toogood AA, Ponder BAJ, Shalet SM. A family with multiple endocrine neoplasia type 2B which does not have a mutation at codon 918 of exon 16 of the *RET* proto-oncogene. J Endocrinol 1995; 144S:P44.
18. Dahia PLM, Aguiar RCT, **Eng C**, Crossey PA, Maher ER, Grossman A, Ponder BAJ, Toledo SPA. Molecular analysis of p53, *VHL*, *RET* genes in pheochromocytoma. The Endocrine Society 1995; 614 (P3-584).
19. Ivanchuk S, **Eng C**, Myers S, Mulligan LM. Expression of alternatively spliced *RET* transcripts in the developing human kidney and Wilms' tumor. Genetics Society of Canada, Guelph, ON, June 1995.
20. Myers SM, **Eng C**, Ponder BAJ, Mulligan LM. Characterization of *RET* proto-oncogene 3' splicing variants and polyadenylation sites: a novel C-terminal for RET. Am J Hum Genet 1995; 37S:A73 (Abstract 389).
21. Ivanchuk S, **Eng C**, Myers S, Mulligan LM. Expression of alternatively spliced *RET* transcripts in the developing human kidney and Wilms' tumor. Am J Hum Genet 1995; 37S:A302 (Abstract 1751).
22. Lin A, Longy M, Goldstein A, Mulvihill J, Ponder B, Tucker M, **Eng C**. Localization of Cowden's disease gene to chromosome 10q22-23. Proc Am Soc Clin Oncol 1996; 15:1746 (Abstract 2003).
23. Mulligan L, **Eng C**, International *RET* Mutation Consortium. *RET* mutations in multiple endocrine neoplasia type 2. International Congress of Endocrinology 1996
24. **Eng C**, Mulligan LM, International *RET* Mutation Consortium. *RET* proto-oncogene mutations in multiple endocrine neoplasia type 2 and medullary thyroid carcinoma. 39e Journées Internationales Klotz d'Endocrinologie Cliniques, 1996
25. Woodward ER, **Eng C**, Affara NA, Ponder BAJ, Maher ER. Molecular genetic investigations of familial pheochromocytoma. J Med Genet 1996; 33 (suppl 1): SP14.

26. **Eng C**, Mulligan LM. The International *RET* Mutation Consortium database: *RET* proto-oncogene mutations in multiple endocrine neoplasia type 2 and related sporadic tumours. 3rd HUGO Mutat Database Mtg, Oct. 29, 1996.
27. **Eng C**, Nelen MR, Marsh DJ, Lin AY, Coulon V, Goldstein AM, Mulvihill JJ, Tucker MA, Zedenius J, Ponder BAJ, Kremer H, Longy M, Padberg GW, International Cowden Consortium. Mapping of the susceptibility gene for Cowden disease to 10q22-23 and genetic analysis in related tumours. Am J Hum Genet 1996; 59S:A6 (Abstract 18).
28. Salomon R, Attié T, Pelet A, Bidaud C, **Eng C**, Munnich A, Lyonnet S. Germline mutations of the Ret ligand, *GDNF*, are not sufficient to cause Hirschsprung disease. Am J Hum Genet 1996; 59S:A283 (Abstract 1640).
29. Eeles RA, Lloyd S, Murday V, Ebbs SR, Davidson J, **Eng C**, Ponder B, Sacks N, Watson M. An evaluation of genetic counselling on risk perception, mental health and cancer-related worry in women at risk of breast cancer: a prospective study in 283 women. Am J Hum Genet 1996; 59S:A334 (Abstract 1949).
30. Mulligan LM, Ivanchuk SM, Campling BG, Sundaresan V, Rabbitts PH, **Eng C**. Analysis of *RET* and *RET* ligand (*GDNF*) in small cell lung carcinoma. Am J Hum Genet 1996; 59S:A345 (Abstract 2010).
31. Nelen MR, von Deimling A, Boerman D, Mariman E, **Eng C**, International Cowden Consortium, Kremer H, Padberg GW. LOH studies in human brain tumors with markers derived from the Cowden critical region. Am J Hum Genet 1996; 59S:A345 (Abstract 2011).
32. Kilgallen C, Smith W, Shapiro S, Cano-Diaz S, Vijg J, **Eng C**, Wolfe HJ. Long chain PCR on archival material for rapid detection of point mutations and molecular screening for prognostic factors in neoplastic progression. Lab Invest 1997; 76:1072
33. Kilgallen C, Diaz-Cano S, **Eng C**, Wolfe HJ. Glial cell line-derived neurotrophic factor (*GDNF*) expression in normal and neoplastic tissues: an immunohistochemical study. Lab Invest 1997; 76
34. Syngal S, Fox E, **Eng C**, Garber JE, Kolodner RD. Presence of more than one *hMSH2* and *hMLH1* mutation in four hereditary nonpolyposis colon cancer (HNPCC) kindreds. Gastroenterol 1997; 112:A664.
35. Schuffenecker I, Virally-Monod, Brohet R, Goldgar D, Conte-Devolx B, Leclerc L, Chabre O, Caron J, Houdent C, Modigliani E, Rohmer V, Schlumberger M, **Eng C**, Guillausseau PJ, Lenoir G, GETC. Risk and penetrance of primary hyperparathyroidism in MEN 2A families with 634 mutations of the *RET* proto-oncogene. Sixth Intl MEN VHL Wrkshp 1997; 78 (Abstract 209).
36. **Eng C**, Mulligan L, International *RET* Mutation Consortium. Genotype-phenotype correlation in MEN 2. Sixth Intl MEN VHL Wrkshp 1997; 98 (Abstract 229).
37. Marsh DJ, Zheng Z, Arnold A, Andrew SD, Learoyd D, Frilling A, Komminoth P, Neumann HPH, Ponder BAJ, Rollins BJ, Shapiro GI, Robinson BG, Mulligan LM, **Eng C**. Mutation analysis of glial cell line-derived neurotrophic factor (*GDNF*), a ligand for the *RET*/co-receptor complex, in MEN 2 and sporadic neuroendocrine tumours. Sixth Intl MEN VHL Wrkshp 1997; 120 (Abstract 251).

38. Woodward ER, Dahia PLM, McMahon R, Voutilainen R, Affara NA, Ponder BAJ, Maher ER, **Eng C**. Genetic predisposition to pheochromocytoma: analysis of candidate genes *GDNF*, *RET* and *VHL*. Sixth Intl MEN VHL Wrkshp 1997: 138 (Abstract 269).
39. Mulligan LM, Timmer T, Ivanchuk SM, **Eng C**, Hofstra RMW. *RET* and its ligand components, *GDNF* and *GDNFR- $\alpha$* , in small cell lung carcinoma. Sixth Intl MEN VHL Wrkshp 1997: 164 (Abstract 323).
40. Marsh DJ, Zheng Z, Zedenius J, Larsson C, Komminoth P, Longy M, **Eng C**. Loss of heterozygosity (LOH) in the region of the Cowden disease locus on 10q22-23 in sporadic thyroid tumours. Sixth Intl MEN VHL Wrkshp 1997: 165 (Abstract 324).
41. Tsou HC, Ping X, Xie X, Yao Y, Schrager C, Gruener A, Christiano AM, Liaw D, Parsons R, **Eng C**, Peacocke M. The molecular basis of Cowden's syndrome. J Invest Dermatol 1997; 109:HB5.
42. **Eng C**, Marsh D, Liaw D, Dahia P, Li J, Zheng Z, Tsou H, Peacocke M, Gorlin R, Parsons R. Germline mutations of the *PTEN* gene in Cowden disease and Bannayan-Zonana syndrome. Am J Hum Genet 1997; 61S:A15 (Abstract 69).
43. Coulon V, Feilotter HE, Boag AH, Dorion-Bonnet F, Duboué B, Latham WCW, **Eng C**, Longy M, Mulligan LM. Loss of heterozygosity of chromosomal region 10q23 in human sporadic breast carcinoma. Am J Hum Genet 1997; 61S:A63 (Abstract 335).
44. Dahia PLM, Marsh DJ, Zheng Z, Zedenius J, Komminoth P, Parsons R, Longy M, Larsson C, **Eng C**. Mutation and deletion analysis of the Cowden disease gene, *PTEN*, in sporadic nonmedullary thyroid tumors. Am J Hum Genet 1997; 61S:A63 (Abstract 338).
45. Feilotter HE, Nagai MA, Boag AH, **Eng C**, Mulligan LM. Analysis of the *PTEN* coding region in primary prostate carcinomas. Am J Hum Genet 1997; 61S:A65 (Abstract 348).
46. Tsou HC, Teng D, Ping XL, Broncolini V, Davis T, Hu R, Xie XX, Gruener AC, Schrager CA, Christiano AC, **Eng C**, Steck P, Ott J, Tavtigian S, Peacocke M. Role of *MMAC1* mutations in early onset breast cancer: causative in association with Cowden syndrome and excluded in *BRCA1*-negative cases. Am J Hum Genet 1997; 61S: A85 (Abstract 468).
47. Myers SM, **Eng C**, Hession C, Cate R, Kogon MD, Mulligan LM. The physical structures of *GDNFR- $\alpha$*  and *NDNR- $\alpha$* . Am J Hum Genet 1997; 61S:A381 (Abstract 2231).
48. Weber HC, Marsh D, Lubensky I, Lin A, **Eng C**. Germline *PTEN/MMAC1/TEP1* mutations and association with gastrointestinal manifestations in Cowden disease. Gastroenterol 1998; 114S: G2902.
49. **Eng C**. Cowden syndrome and Bannayan-Ruvalcaba-Riley syndrome: *PTEN* mutation analysis as a molecular diagnostic tool? Horm Metab Res 1998; 30:288-9.
50. **Eng C**. Cowden syndrome: update on genetic mechanisms and clinical features. J Cancer Res Clin Oncol 1998; 124: S16 (TL14).

51. **Eng C**, Mulligan LM, International *RET* Mutation Consortium. Genotype-phenotype correlations in MEN 2 and genotype-prognosis studies in sporadic medullary thyroid carcinoma. J Invest Endocrinol (in press)
52. **Eng C**, Smith WM, van Orsouw NJ, Dhanda RK, Rines RD, Sigalas I, Fox EA, Kolodner RD, Vijg J. DGGE-based two-dimensional gene scanning as a rapid, accurate, cost-effective technology for snapshot detection of mutations. Am J Hum Genet 1998; 63S:A20 (Abstract 145)
53. Gimm O, Neuberg DS, Marsh DJ, Dahia PLM, Hoang-Vu C, Raue F, Hinze R, Dralle H, **Eng C**. Over-representation of a germline *RET* sequence variant in patients with sporadic medullary thyroid carcinoma and somatic *RET* codon 918 mutation. Am J Hum Genet 1998; 63S:A20 (Abstract 102)
54. Dabora S, Sigalas I, Peters S, **Eng C**, Vijg J, Kwiatkowski D. Comprehensive mutational analysis of *TSC1* by two-dimensional DGGE. Am J Hum Genet 1998; 63S:A229 (Abstract 1313)
55. Gössling A, Gimm O, Marsh DJ, Dahia PLM, Myers SM, Mulligan LM, von Deimling A, **Eng C**. Somatic deletion of the gene encoding GFR $\alpha$ -1, a co-receptor of RET, in sporadic brain tumours. Am J Hum Genet 1998; 63S:A71 (Abstract 377)
56. Cummings S, Marsh DJ, Sveen LW, **Eng C**, Olopade OI. Cowden syndrome (CS) in African American (AA) kindreds: identification of germline *PTEN* mutation in a male breast cancer patient. Am J Hum Genet 1998; 63S:A66 (Abstract 349)
57. Danziger K, Syngal S, Fox E, **Eng C**, Kolodner R, Garber J. *hMLH1* and *hMSH2* mutations in patients with early-onset colorectal cancer (CRC). Am J Hum Genet 1998; 63S:A66 (Abstract 350)
58. Maher ER, Clifford SC, Walsh S, Hewson K, Brinke A, Green PM, Gianelli F, **Eng C**. Characterisation and mutation analysis of candidate renal cell carcinoma genes (Cul2 and VBP-1). Am J Hum Genet 1998; 63S:A77 (Abstract 414)
59. Dasouki MJ, Roe CR, Butler MG, **Eng C**. Fatty myopathy in Bannayan-Zonana syndrome is not due to LCHAD (long chain 3-hydroxy acyl co-A dehydrogenase) deficiency. Am J Hum Genet 1998; 63S:A265 (Abstract 1526)
60. Marsh DJ, Kum JB, Lunetta KL, Bennet MJ, Hunter A, Gorlin RJ, Dahia PLM, **Eng C**. Germline mutation analysis and genotype-phenotype correlations in Bannayan-Riley-Ruvalcaba syndrome. Proc Am Assoc Cancer Res 1999; 40:272 (Abstract 1804)
61. Dahia PLM, Aguiar RCT, Alberta J, Kum JB, Caron S, Heinz S, Marsh DJ, Ritz J, Freedman A, Stiles C, **Eng C**. PTEN is inversely correlated with the cell survival factor Akt/PKB and is inactivated via multiple mechanisms in haematologic malignancies. Proc Am Assoc Cancer Res 1999; 40:281-2 (Abstract 1869)
62. **Eng C**. The role of *PTEN* in Cowden syndrome and multiple sporadic cancers. Proc Seventh Intl MEN Wkshop 1999; 147-52 (Plenary)
63. Gimm O, Greco A, Hoang-Vu C, Dralle H, Pierotti MA, **Eng C**. Mutation analysis reveals novel sequence variants in *NTRK1* in sporadic human medullary thyroid carcinoma. Proc Seventh Intl MEN Wkshop 1999; 265 (Poster Th 22)

64. Gimm O, Neuberg DS, Marsh DJ, Dahia PLM, Hoang-Vu C, Raue F, Hinze R, Dralle H, **Eng C**. Over-representation of the germline *RET* sequence variant S836S in patients with sporadic medullary thyroid carcinoma and somatic *RET* codon 918 mutation. Proc Seventh Intl MEN Wkshop 1999; 266 (Poster Th 23)
65. Tisell L-E, Nilsson O, Jansson S, Nilsson B, Ahlman H, Gimm O, **Eng C**. Adrenal and extra-adrenal pheochromocytomas in a family with germline *RET* V804L mutations, previously associated only with familial medullary thyroid carcinoma. Proc Seventh Intl MEN Wkshop 1999; 282 (Poster Th 37)
66. Gimm O, Gössling A, Marsh DJ, Dahia PLM, Mulligan LM, von Deimlin A, **Eng C**. Mutation and deletion analysis of GFR $\alpha$ -1, encoding the coreceptor for the GDNF/RET complex, in human brain tumours. Proc Seventh Intl MEN Wkshop 1999; 284 (Poster Th 38)
67. **Eng C**, Sarraf P, Mueller E, Smith WM, Wright HM, Kum JB, Aaltonen LA, de la Chapelle A, Spiegelman BM. Loss of function mutations in PPARgamma in sporadic primary colorectal cancer. Am J Hum Genet 1999; 64S (in press)
68. Duerr EM, Gimm O, Kum JB, Clifford SC, Toledo SPA, Maher ER, Dahia PLM, **Eng C**. Two polymorphisms in the VHL-associated gene *CUL2* are over-represented in pheochromocytoma patients without somatic *CUL2* mutations. Am J Hum Genet 1999; 64S (in press)
69. Gimm O, Borrego S, Saez ME, Ruiz A, López-Alonzo M, **Eng C**, Antiñolo G. Specific sequence polymorphisms in the *RET* proto-oncogene are over-represented in individuals with Hirschsprung disease and may represent loci modifying phenotypic expression. Am J Hum Genet 1999; 64S (in press)
70. Yeh JJ, Lunetta KL, Dahia PLM, **Eng C**. Mitochondrial DNA (mtDNA) mutations in papillary thyroid carcinoma and differential mtDNA sequence variants in cases with malignant versus benign thyroid tumors. Am J Hum Genet 1999; 64S (in press)
71. Hampel H, Poling BA, Curtis M, Mascari M, Fromkes JJ, **Eng C**. Familial Barrett esophagus: a true hereditary cancer syndrome. Am J Hum Genet 1999; 64S (in press)

#### Newsletters and Misc.:

1. Eng C, Li FP. Retinoblastoma in families. New England Retinoblastoma Support Group, Spring 1991.
2. Writer for Trends in Genetics Monitor, 1994